

RESEARCH ARTICLE

Open Access

The *Aspergillus giganteus* antifungal protein AFP_{NN5353} activates the cell wall integrity pathway and perturbs calcium homeostasis

Ulrike Binder^{1,5}, Mojca Bencina^{2,3}, Andrea Eigentler¹, Vera Meyer⁴ and Florentine Marx^{1*}

Abstract

Background: The antifungal protein AFP_{NN5353} is a defensin-like protein of *Aspergillus giganteus*. It belongs to a group of secretory proteins with low molecular mass, cationic character and a high content of cysteine residues. The protein inhibits the germination and growth of filamentous ascomycetes, including important human and plant pathogens and the model organisms *Aspergillus nidulans* and *Aspergillus niger*.

Results: We determined an AFP_{NN5353} hypersensitive phenotype of non-functional *A. nidulans* mutants in the protein kinase C (Pkc)/mitogen-activated protein kinase (Mpk) signalling pathway and the induction of the α -glucan synthase A (*agsA*) promoter in a transgenic *A. niger* strain which point at the activation of the cell wall integrity pathway (CWIP) and the remodelling of the cell wall in response to AFP_{NN5353}. The activation of the CWIP by AFP_{NN5353}, however, operates independently from RhoA which is the central regulator of CWIP signal transduction in fungi.

Furthermore, we provide evidence that calcium (Ca^{2+}) signalling plays an important role in the mechanistic function of this antifungal protein. AFP_{NN5353} increased about 2-fold the cytosolic free Ca^{2+} ($[\text{Ca}^{2+}]_c$) of a transgenic *A. niger* strain expressing codon optimized aequorin. Supplementation of the growth medium with CaCl_2 counteracted AFP_{NN5353} toxicity, ameliorated the perturbation of the $[\text{Ca}^{2+}]_c$ resting level and prevented protein uptake into *Aspergillus sp.* cells.

Conclusions: The present study contributes new insights into the molecular mechanisms of action of the *A. giganteus* antifungal protein AFP_{NN5353}. We identified its antifungal activity, initiated the investigation of pathways that determine protein toxicity, namely the CWIP and the Ca^{2+} signalling cascade, and studied in detail the cellular uptake mechanism in sensitive fungi. This knowledge contributes to define new potential targets for the development of novel antifungal strategies to prevent and combat infections of filamentous fungi which have severe negative impact in medicine and agriculture.

Background

All organisms have evolved several defence systems in order to protect themselves against bacteria, fungi and viruses. Higher organisms have developed a complex network of humoral and cellular responses, called adaptive immunity. A second defence system, the innate immunity, consists of many components, including small peptides with a broad antimicrobial spectrum [1,2]. The production of such proteins with

antimicrobial activity is not limited to higher eukaryotes, but also found in microorganisms, including fungi. The diversity of these proteins is reflected in their mode of action and their species-specificity. Some of them form pores in the membrane, others are known to inhibit cell wall synthesis or interfere with nucleic acids and their synthesis [3,4]. They can be involved in the inhibition of protein synthesis or interfere with cell cycle control [3,4]. A relatively new group of antimicrobial proteins secreted by filamentous ascomycetes includes small, cationic and cysteine-rich proteins. So far, only few antifungal proteins have been characterized, namely AFP from *Aspergillus giganteus*, ANAFP from *Aspergillus*

* Correspondence: florentine.marx@i-med.ac.at

¹Biocenter, Division of Molecular Biology, Innsbruck Medical University, Fritz-Pregl Strasse 3, Innsbruck, A-6020, Austria

Full list of author information is available at the end of the article

niger, PAF from *Penicillium chrysogenum* and NAF from *Penicillium nalgiovense* [5-8].

The mode of action of these proteins is not fully understood. Nevertheless, there is evidence, that their toxicity is mediated by interaction with distinct molecules or receptors at the outer layers of the cell, e.g. cell wall or plasma membrane. Deleterious effects can then be induced either by transmitting signals from the outer layers into the cell, or by internalization of the protein and interaction with internal molecules [9-15]. Similar to substances that perturb the cell wall, such as caspofungin, congo red or calcofluor white (CFW) [10,16], the *A. giganteus* antifungal protein AFP was found to modulate the cell wall composition by enhancing the expression of the α -1,3-glucan synthase A gene (*agsA*), possibly by the activation of the cell wall integrity pathway (CWIP), and inhibiting chitin synthesis in sensitive fungi [10]. This, however, stands in contrast to the mode of action of the *P. chrysogenum* antifungal protein PAF which fails to activate the CWIP [9]. However, the central players that trigger cell wall remodelling in AFP-sensitive fungi have not been investigated so far.

Another mechanistic function of antifungal proteins is the interference with ion, especially Ca^{2+} ion homeostasis and signalling [15,17,18]. We could recently show that the *P. chrysogenum* antifungal protein PAF severely perturbed the Ca^{2+} homeostasis of *Neurospora crassa* by rapidly elevating the cytoplasmic Ca^{2+} [Ca^{2+}]_c resting level [17]. Numerous reports indicate that the activity of antifungal proteins can be antagonized by the external addition of Ca^{2+} ions to the test medium [15,17-21] pointing towards the induction of adaptive responses which may be triggered by Ca^{2+} signalling [15,17].

The aim of this study was to characterize in more detail the mode of action of the *A. giganteus* AFP variant protein AFP_{NN5353} and to investigate the pathways that might be affected/modulated by this antifungal protein. Therefore, we focussed our interest on the involvement of the CWIP and the Ca^{2+} signalling in the toxicity of AFP_{NN5353}. To address these questions, we used the highly AFP_{NN5353} sensitive model organisms *A. nidulans* and *A. niger* for which appropriate mutant strains were available.

Results

In silico analysis of AFP_{NN5353}

CLUSTALW amino acid (aa) sequence analysis of AFP_{NN5353} with other known antifungal proteins revealed that AFP_{NN5353} from *A. giganteus* strain A3274 is a protein homologous to AFP from *A. giganteus* strain MDH 18894 [8,22]. AFP_{NN5353} exhibits > 90% identity with AFP, but only 42% identity with the *P. chrysogenum* PAF and 27% identity with the *A. niger* ANAFP. In fact, the secreted mature form of AFP_{NN5353} consists

of 51 aa and differs only in 5 aa from AFP (Figure 1). Three aa exchanges belong to structurally related aa, one aa exhibits weak similarity and one aa is different (position 4). These aa exchanges do not influence the theoretical isoelectric point (pI) of AFP_{NN5353}, which is the same as for AFP (pI 9.3, <http://expasy.org/tools/protparam.html>). Most importantly, AFP_{NN5353} still contains the putative chitin-binding domain CKYKAQ present in AFP but not in PAF or ANAFP and also harbors all conserved cysteine residues important for protein stabilization [10,23].

Antifungal activity of the protein AFP_{NN5353}

To investigate the antifungal specificity of AFP_{NN5353}, fifteen filamentous fungi were tested for their susceptibility to the protein. Since antifungal proteins might be useful for biotechnological applications, filamentous human and plant pathogenic fungi were selected as test organisms (e.g. *Fusarium oxysporum*, *Botrytis cinerea*, *Mucor* sp. and *A. fumigatus*) in addition to the model organisms *A. nidulans* and *A. niger*. As shown in Table 1, thirteen out of fifteen tested moulds were found to be sensitive against AFP_{NN5353}. *A. nidulans* wild type, *N. crassa* wild type and *A. niger* wild type were the most sensitive strains to AFP_{NN5353}. The minimal inhibitory concentration (MIC) of AFP (the concentration that completely inhibited conidial germination in liquid growth assays) was 0.2 $\mu\text{g/ml}$ for *A. nidulans*, 0.5 $\mu\text{g/ml}$ for *N. crassa* and 1 $\mu\text{g/ml}$ for *A. niger*. Two strains were unaffected at the protein concentrations tested: *M. circenelloides* and *M. genevensis* were insensitive against AFP_{NN5353} when concentrations up to 500 $\mu\text{g/ml}$ were used.

AFP_{NN5353} interferes with the cell wall integrity of *A. nidulans*

It is known that antifungal compounds such as congo red, caffeine, CFW or caspofungin interfere with cell wall biosynthesis and weaken the cell wall in fungi (reviewed by [24]). The remodeling of the cell wall by these antifungal compounds is mediated by the activation of the CWIP. In fungi, extracellular signals are transmitted via the membrane bound small GTPase RhoA to the central regulators Pkc and Mpk, which are regulated by phosphorylation/dephosphorylation. The signal transduction cascade eventually enforces transcription of cell wall synthesis genes, partly via the transcription factor RlmA [16,25]. Respective loss-of-function or conditional mutants show hypersensitive phenotypes in the presence of cell wall perturbing agents [9,24-26]. Similar to substances that weaken the cell wall, the *A. giganteus* antifungal protein AFP modulates the cell wall composition by inhibiting chitin synthesis in sensitive fungi (e.g. *A. niger*, *A. oryzae*) and

```

AFPNN5353 -ATYPGKCYKKDNICKYKAQSGKTGICKC---YVKRCPRDGAKCELDSYKGGKCHC----
AFP      -ATYNGKCYKKDNICKYKAQSGKTAICKC---YVKKCPRDGAKCEFDSYKGGKCYC----
ANAFP   LSKYGGECSLEHNTCTYRKD-GKNHVVSCPSAANLRCKTDRHHCEYDDHHKTVDCQTPV
PAF     -AKYTGKCTKSKNECKYKNDAGKDTFIKCPKFDNKKCTKDNNKCTVDTYNNAVDCD---
          :.* *:* ..* *.* ** . .* :* : *:* * :.*
    
```

Figure 1 Clustalw sequence alignment <http://www.ebi.ac.uk/Tools/msa/clustalw2/> of the antifungal proteins AFP_{NN5353} and AFP from *A. giganteus*, ANAFP from *A. niger* and PAF from *P. chrysogenum*. Identical amino acids (aa) are marked with (*), aa with strong similarity are indicated with (:) and aa with weak similarity are marked with (.)

inducing the expression of *agsA* most likely by the activation of the CWIP [10].

To study the involvement of the CWIP in AFP_{NN5353} toxicity, we first tested whether the osmotic stabilizer sorbitol counteracts the toxicity of AFP_{NN5353}. In the absence of AFP_{NN5353} *A. nidulans* proliferated less well in the presence of 1 M sorbitol and reached only 30% growth compared to the growth in standard medium (100%). Nevertheless, the addition of 1 M sorbitol to the growth medium strongly reduced the activity of AFP_{NN5353} on *A. nidulans* wild type. The osmotic stabilizer ameliorated growth in the presence of 0.05 µg/ml AFP_{NN5353} by 80% compared to a 10% growth rate in the absence of sorbitol (Table 2). This was even more accentuated when 0.1 and 0.2 µg/ml AFP_{NN5353} were applied, suggesting that AFP_{NN5353} indeed weakens the cell wall of *A. nidulans*.

To investigate whether AFP_{NN5353} induces *agsA* gene transcription similar to AFP via the Pkc/Mpk signalling pathway, we tested the effect of the antifungal protein on the transgenic *A. niger* strain RD6.47 which expresses

a nuclear-targeted GFP protein fused to the *A. niger agsA* promoter. RD6.47 germlings were treated with AFP_{NN5353} (conc. 10 to 100 µg/ml) for 2 h and analyzed microscopically. As shown in Additional file 1, a nuclear signal was clearly detectable in germlings of RD6.47 treated with ≥ 50 µg/ml AFP_{NN5353}, similar to that when exposed to 10 µg/ml caspofungin. In untreated germlings, however, no signal could be observed. These observations perfectly match with the data obtained for AFP [10]. It has to be noted here that antifungal protein concentrations higher than the MIC determined for conidia (> 10-50 fold) are needed to inhibit the growth of germlings or hyphae of sensitive fungi [10,27] (data not shown).

Next, we tested several *A. nidulans* mutant strains affected in central players of the CWIP for their susceptibility to AFP_{NN5353} by determining their radial growth in the presence or absence of the antifungal protein. Since RhoA is an essential protein in *A. nidulans*, two strains with ectopic copies of the constitutively active *rhoA*^{G14V} allele and the dominant *rhoA*^{E40I} allele [28] were tested in comparison to the wild type strain (GR5). The *rhoA*^{G14V} mutation prevents the hydrolysis of GTP and therefore renders RhoA constantly active [28]. Similarly, the GTP hydrolysis is inhibited in the *RhoA*^{E40I} strain, but this mutation also perturbs the binding of the GTPase activating protein (GAP) to RhoA and possibly disturbs downstream effectors of RhoA-GAP [28]. The constitutively active *RhoA*^{G14V} and the dominant *RhoA*^{E40I} strain exhibited the same sensitivity towards AFP_{NN5353} as the wild type strain at low protein concentrations (≤ 0.2 µg/ml) (Figure 2A). Interestingly, the dominant *RhoA*^{E40I} strain was more

Table 1 Minimal inhibitory concentrations (MIC; µg/ml) of AFP_{NN5353} against different filamentous fungi


organism	MIC (µg/ml)
<i>Aspergillus flavus</i> ATCC9643	50
<i>Aspergillus fumigatus</i> ATCC 46645	50
<i>Aspergillus giganteus</i> AG090701	50
<i>Aspergillus nidulans</i> FGSC4	0.2
<i>Aspergillus niger</i> CBS 120.49	1
<i>Aspergillus terreus</i> 304	5
<i>Botrytis cinerea</i> BC 080801	10
<i>Fusarium oxysporum</i> FO 240901	5
<i>Fusarium sambucinum</i> FS210901	5
<i>Gliocladium roseum</i> GR 210901	100
<i>Mucor circinelloides</i> MC080801	insensitive ^a
<i>Mucor genevensis</i> MG 080801	insensitive ^a
<i>Penicillium chrysogenum</i> ATCC10002	10
<i>Trichoderma koningii</i> TC 060901	20
<i>Neurospora crassa</i> FGSC 2489	0.5

^aup to 500 µg/ml AFP_{NN5353} was tested
 1 × 10⁴ conidia/ml were incubated in 200 µl CM medium in the presence of various concentrations of AFP_{NN5353} at 30°C for 24 h. Growth was determined by measuring the OD_{620 nm}.

Table 2 The effect of 1 M sorbitol on the growth inhibiting activity of AFP_{NN5353} on *A. nidulans*

AFP _{NN5353} (µg/ml)	CM	CM + 1 M sorbitol
0	100 (SD ± 10)	100 (SD ± 11)
0.05	10.4 (SD ± 1)	79.3 (SD ± 6)
0.1	5.5 (SD ± 2)	68.3 (SD ± 0.8)
0.2	no growth	17.8 (SD ± 0.8)

1 × 10⁴ conidia/ml were incubated in CM with 0-0.2 µg/ml AFP_{NN5353} for 24 h. Percent values were calculated from percent changes in OD₆₂₀ of AFP_{NN5353} treated *A. nidulans* compared to untreated controls (= 100%). Results are expressed as mean ± SD (n = 3).



resistant to AFP_{NN5353} than the wild type strain or the RhoA^{G14V} strain at higher protein concentrations (1 μg/ml) (Figure 2A). Therefore, we suggest that the toxicity of AFP_{NN5353} is transmitted by RhoA-GAP targets and not by RhoA itself. These mutants performed similarly when exposed to the orthologous *P. chrysogenum* anti-fungal protein PAF [9].

In addition, mutants defective in PkcA and MpkA activity were tested for their AFP_{NN5353} susceptibility. As *pkcA* is an essential gene in *A. nidulans*, a conditional *alcA*-PKC mutant strain was used, where the *pkcA* gene was put under the control of the *alcA*

promoter, which is repressed by glucose but derepressed by glycerol [26]. Both the conditional *alcA*-PKC mutant (cultivated under repressive conditions) and a *ΔmpkA* mutant were hypersensitive to AFP_{NN5353} compared to their recipient strains R153 and GR5, respectively, indicating that the activity of PkcA and MpkA confers a certain resistance to AFP_{NN5353} (Figure 2A). The hypersensitive phenotype of the *ΔmpkA* mutant was also confirmed by liquid growth inhibitory assays. In unchallenged liquid condition, the GR5 and the *ΔmpkA* mutant showed a comparable proliferation rate (Figure 2B). In the presence of 0.05 μg/ml AFP_{NN5353}, however,

the *mpkA* deletion strain did not germinate whereas the GR5 strain still exhibited 11% growth. Note that growth inhibition in liquid conditions requires less antifungal protein to monitor its toxicity than on solid media probably due to less diffusion in the latter case (data not shown).

From these data we conclude that AFP_{NN5353} interferes with the cell wall homeostasis of *A. nidulans* and that this interaction is mediated by PkcA/MpkA signaling, although independently from RhoA.

AFP_{NN5353} disrupts calcium homeostasis in *A. niger*

Supplements other than osmotic stabilizers can also antagonize the activity of antifungal proteins from plants and ascomycetes. For example, the addition of cations such as Ca²⁺ ions to the growth medium reversed the antifungal activity of the *P. chrysogenum* PAF [17], the *A. giganteus* AFP [15,21] and of plant defensins [29,30] which are usually positively charged due to their high pI. A cation-sensitive antifungal mode of action can for example be associated with the perturbation of the intracellular Ca²⁺ homeostasis by antifungal peptides [17,18] but might also result from the interference of cations with antifungal-target interaction(s).

Therefore, we tested to which extent these effects also account for the antifungal activity of AFP_{NN5353}. To this end, we selected *A. niger* as model organism because this mould was highly sensitive to AFP_{NN5353} and a transgenic strain was available that expressed the recombinant codon optimized Ca²⁺-sensitive photoprotein aequorin for measuring the [Ca²⁺]_c resting level in response to AFP_{NN5353} [31]. First, we tested the activity of AFP_{NN5353} in Vogels* medium supplemented with 5-20 mM CaCl₂ or without CaCl₂ as a control (data not shown). Addition of CaCl₂ did not influence the growth of *A. niger* up to a concentration of 20 mM. The growth of *A. niger* exposed to AFP_{NN5353}, however, ameliorated in the presence of increasing concentrations of CaCl₂. 20 mM CaCl₂ neutralized the toxicity of 0.5-1.0 µg/ml AFP_{NN5353} and the treated samples resumed growth to 100% (Table 3).

Next, we determined the influence of AFP_{NN5353} on the intracellular Ca²⁺ signature. Before AFP_{NN5353}

Table 3 The effect of 20 mM external CaCl₂ (in Vogels* medium) on the growth inhibitory activity of AFP_{NN5353} on *A. niger* strain A533.

AFP _{NN5353} (µg/ml)	Vogels*	Vogels* + 20 mM Ca ²⁺
0	100 (SD ± 10)	100 (SD ± 8)
0.5	12 (SD ± 3)	101 (SD ± 9)
1.0	no growth	105 (SD ± 6)

OD₆₂₀ was measured after 24 h of incubation. The growth of untreated controls was normalized to 100% to evaluate the percent growth of samples in the presence of AFP_{NN5353}. Vogels* medium without CaCl₂ supplementation contains 0.7 mM Ca²⁺. Results are expressed as mean ± SD (n = 3).

addition, the resting level of the intracellular Ca²⁺ was 0.08 µM. We could show, however, that the [Ca²⁺]_c resting level was significantly increased in twelve h old *A. niger* cultures that were treated with 20 µg/ml AFP_{NN5353}. The [Ca²⁺]_c resting level rose to a maximum of 0.19 µM within the first 8 min and stayed elevated throughout the time of measurement (60 min), whereas the Ca²⁺ level of the untreated control remained at 0.08 µM (Figure 3). This indicated that AFP_{NN5353} indeed disrupts Ca²⁺ homeostasis in *A. niger*.

To exclude the possibility that the AFP_{NN5353} induced rise in the [Ca²⁺]_c resting level is due to membrane permeabilization and/or pore formation, we studied the effects of AFP_{NN5353} on germlings in the presence of CMFDA, a membrane permeant dye that is metabolized by viable cells, and the membrane impermeant dye propidium iodide (PI). Additional file 2 shows that samples treated with 20 µg/ml AFP_{NN5353} for 10 min metabolized CMFDA but did not take up PI, resulting in green but no red fluorescence, similar to untreated controls. This indicated that the plasma membrane was still intact after 10 min of protein treatment. Samples exposed to ethanol did not metabolize CMFDA but appeared bright red due to PI internalization, indicating that here the membrane was permeabilized. We therefore conclude that the rapid increase in [Ca²⁺]_c within the first 10 min of protein treatment is not the result of uncontrolled Ca²⁺ influx due to plasma membrane permeabilization.

The calcium chelator BAPTA abrogates the AFP_{NN5353}-induced calcium signature

The increased [Ca²⁺]_c in response to AFP_{NN5353} treatment could originate from extracellular and/or from intracellular Ca²⁺ stores, such as mitochondria, vacuoles,




Figure 3 Increase in resting [Ca²⁺]_c of twelve h old *A. niger* germlings treated with AFP_{NN5353} or no protein (controls). Measurements were taken every 1.4 minutes. Values represent average of six samples.

endoplasmic reticulum or the Golgi apparatus. To discriminate between the extracellular and intracellular source of the $[Ca^{2+}]_c$ increase, we tested the influence of the Ca^{2+} -selective membrane impermeable chelator BAPTA. On its own, BAPTA did not influence the resting level of $[Ca^{2+}]_c$ in twelve h old *A. niger* cultures (Figure 4). However, a pretreatment of the samples with 10 mM BAPTA before the addition of AFP_{NN5353} inhibited the protein-specific increase in $[Ca^{2+}]_c$ resting level (Figure 4). Interestingly, the elevated $[Ca^{2+}]_c$ in response to a 40 min AFP_{NN5353}-treatment dropped to the resting level immediately after the addition of 10 mM BAPTA (Figure 4), indicating that the AFP_{NN5353}-induced elevation of the $[Ca^{2+}]_c$ resting level requires the continuous influx of extracellular Ca^{2+} and eventually results in loss of $[Ca^{2+}]_c$ homeostasis.

Extracellular calcium ameliorates the AFP_{NN5353}-induced rise in $[Ca^{2+}]_c$

To decipher the observation that high external $CaCl_2$ concentrations counteracted AFP_{NN5353} toxicity (Table 3), we monitored the effect of externally added Ca^{2+} on the AFP_{NN5353}-induced Ca^{2+} signature. To this end, *A. niger* germlings were preincubated with 20 mM $CaCl_2$ for 10 min before 20 μ g/ml AFP_{NN5353} was added and the changes in the $[Ca^{2+}]_c$ resting level were monitored over a time course of 60 min. This treatment resulted in a less pronounced rise of the $[Ca^{2+}]_c$ resting level compared to samples without preincubation with $CaCl_2$. In contrast, the presence of 20 mM $CaCl_2$ alone had no major effect on the intracellular $[Ca^{2+}]_c$ resting level which resembled that of the control without AFP_{NN5353} (data not shown). The values of the $[Ca^{2+}]_c$ resting levels of the last 10 min (50 to 60 min) measurement of




Figure 4 Effect of the extracellular chelator BAPTA on the AFP_{NN5353} induced $[Ca^{2+}]_c$ resting level. 10 mM BAPTA (final conc.) were applied 40 min before or 40 min after treatment with 20 μ g/ml AFP_{NN5353}. Samples without supplements were used as controls. SD (n = 6) was less than 10% of the values presented.

AFP_{NN5353} treatment in the presence or absence of high Ca^{2+} concentration (20 mM versus 0.7 mM) are summarized in Table 4. The average of the $[Ca^{2+}]_c$ of the controls which were not exposed to AFP_{NN5353} was 0.039 μ M in the presence of 0.7 μ M $CaCl_2$ (standard condition) and 0.062 μ M in the presence of 20 mM $CaCl_2$. When AFP_{NN5353} was added, there was no significant elevation of the $[Ca^{2+}]_c$ in high- Ca^{2+} medium (20 mM) (0.057 μ M) whereas the $[Ca^{2+}]_c$ rised to 0.146 μ M at standard $CaCl_2$ concentration (0.7 mM). These results suggest that Ca^{2+} externally added prior to the addition of AFP_{NN5353} counteracts the AFP_{NN5353} induced perturbation of the $[Ca^{2+}]_c$ and growth inhibitory effect, at least partly, by controlling the $[Ca^{2+}]_c$ resting level.

AFP_{NN5353} decreases the amplitude of the $[Ca^{2+}]_c$ response to mechanical perturbation in *A. niger*

It is known that a range of external stimuli transiently increase $[Ca^{2+}]_c$ levels in *Aspergilli* and other fungi [31,32]. One of these physiological stimuli is mechanical perturbation, which is achieved by the rapid injection of isotonic medium into the test system. This stimulus results in a unique Ca^{2+} signature, likely involving different components of the Ca^{2+} -signalling and Ca^{2+} homeostatic machinery. Changes in this specific Ca^{2+} signature in the presence of compounds, such as AFP_{NN5353}, can give insights into the mode of action of these compounds. In our study, twelve h old cultures of *A. niger* were pre-incubated with AFP_{NN5353} for 60 min and thereafter subjected to mechanical perturbation (rapid injection of 100 μ l Vogels medium). The resulting Ca^{2+} signature, including $[Ca^{2+}]_c$ resting level, kinetics and amplitude, were determined and compared with controls that were not exposed to the protein but also subjected to mechanical perturbation. As shown in Figure 5, AFP_{NN5353} provoked a less pronounced $[Ca^{2+}]_c$ amplitude; however, the $[Ca^{2+}]_c$ level remained elevated even after the stimulus specific response had stopped.

AFP_{NN5353} binding and uptake are essential for protein toxicity in *A. nidulans*

To understand the function of antifungal proteins, the identification of the site of action in target organisms is crucial. So far, controversial reports exist of the

Table 4 The effect of high external $CaCl_2$ concentration on the AFP_{NN5353} induced Ca^{2+} signature in response to AFP_{NN5353}.

$[CaCl_2]$ in Vogels*	0 μ g/ml AFP _{NN5353}	20 μ g/ml AFP _{NN5353}
0.7 mM	0.039 (SD \pm 0.001)	0.146 (SD \pm 0.009)
20 mM	0.062 (SD \pm 0.003)	0.057 (SD \pm 0.004)

Twelve h old germlings were preincubated with 20 mM $CaCl_2$ for 10 min before exposure to AFP_{NN5353}. Values represent the average μ M concentration of $[Ca^{2+}]_c$ within the last 10 min (50-60 min) of measurement.




Figure 5 Effects of AFP_{NN5353} on the [Ca²⁺]_c response to mechanical perturbation. Twelve h old *A. niger* cultures were treated with 20 µg/ml AFP_{NN5353} for 60 min before stimulation by mechanical perturbation (addition of 100 µl Vogels medium). The [Ca²⁺]_c signature was monitored for 5 min. Values represent the average of six samples.

localization of the homologous *A. giganteus* AFP protein. AFP has been detected to bind to outer layers, e.g. the cell wall or the plasma membrane of sensitive fungi [20,21] and a time- and concentration-dependent intracellular localization was reported [20]. In another study, Alexa-labelled AFP was shown to be internalized by the fungal cell and to localize to the nucleus [33].

To dissect the uptake and localization of AFP_{NN5353}, we performed indirect immunofluorescence staining with *A. nidulans* wild type exposed to a sublethal concentration of AFP_{NN5353} (0.2 µg/ml). We applied a protein amount below the toxic concentration for hyphae to maintain the cellular structure and to avoid apoptotic cell disruption [34]. Our study revealed that the protein was internalized after 90 min of incubation, mostly in hyphal tips, but also within hyphal segments (Figure 6A, B). The protein seemed not to localize to cell compartments, but was distributed in the cytoplasm. Similar results were obtained with *A. niger* wild type (data not shown). Control experiments proved the specificity of the intracellular immunofluorescent signals: no intracellular fluorescent signals were detected in samples where either AFP_{NN5353} (Figure 6C, D) or the primary antibody or the secondary antibody was omitted (data not shown).

To analyse the AFP_{NN5353} localization in more detail, *A. nidulans* was incubated with AFP_{NN5353} in the presence of latrunculin B, a potent inhibitor of actin


polymerization and endocytosis [35-37]. At low latrunculin B concentrations (5 µg/ml), protein uptake was severely reduced compared to the positive control without latrunculin B (data not shown), whereas 20 µg latrunculin B/ml completely inhibited the uptake of 0.2 µg/ml AFP_{NN5353}. The solvent of latrunculin B, DMSO, had no adverse effect on protein uptake (data not shown). This indicates that AFP_{NN5353} enters the *A. nidulans* cells by an endocytotic mechanism (Figure 6E, F).

Based on our observation that Ca²⁺ ions antagonize the growth inhibitory activity of AFP_{NN5353}, we questioned whether Ca²⁺ prevents actin-mediated internalisation of the antifungal protein. Indeed, the presence of 10 mM CaCl₂ inhibited protein uptake (Figure 6G, H). Most interestingly, no specific fluorescent signals were detectable in *M. circinelloides* when treated with up to 500 µg/ml of antifungal protein (data not shown), indicating that AFP_{NN5353} does not bind to insensitive strains.

Discussion

In this study we provide important insights into the mechanistic basis of AFP_{NN5353}, a AFP homologous protein.

Species specificity tests revealed that AFP_{NN5353} is active against a broad range of filamentous fungi, including human and plant pathogens. Although the



proteins AFP_{NN5353} and AFP are almost identical and show a similar toxicity, MICs for AFP_{NN5353} differed slightly from those reported for AFP [21]. We attribute this discrepancy to differences in the experimental setups, e.g. fungal strains, medium composition, conidial inoculum, incubation times, cultivation temperature etc., rather than to the differences in the primary sequence of both proteins.

It has been reported that the closely related AFP protein interfered with cell wall synthesis [10] and our finding that the osmotic stabilizer sorbitol neutralized AFP_{NN5353} toxicity further corroborated this assumption. Two *A. nidulans* mutants, the conditional *alcA*-PkcA and the *mpkA* deletion mutant showed a hypersensitive phenotype when exposed to AFP_{NN5353}. This is in

agreement to the reported function of cell wall stressing agents, such as CFW or caffeine in *S. cerevisiae* and *A. nidulans* [9,16,24,26,38,39] and to the *Penicillium* antifungal protein PAF [9]. Importantly, Mpk function is essential for CWIP activation in both, unicellular and filamentous fungi [10,16,40] and triggers the activation of the transcription factors Rlm1p and SBF which regulate the expression of cell cycle regulated genes and genes involved in the synthesis and remodelling of the fungal cell wall in *S. cerevisiae* [41,42]. Similarly, RlmA dependent induction of the expression of the *ags* gene was also reported for aspergilli [25]. Importantly, the activation of the CWIP can occur in a RhoA-dependent, e.g. with CFW [9,43], or RhoA-independent way, the latter proved for PAF and caffeine [9,16] and for AFP_{NN5353} (this study). As proposed by [28] the dominant *rhoA*^{E40I} allele suffers from a perturbation of its GAP binding domain and downstream effectors of Rho-GAP might be disturbed. Therefore, we hypothesize that Rho-GAP targets might be involved in the toxicity of AFP_{NN5353} similarly to the mode of action of the *P. chrysogenum* PAF [9]. Our assumption of the activation of the CWIP by AFP_{NN5353} was further strengthened by the fact, that AFP_{NN5353} treatment induced *agsA* expression in the *A. niger* reporter strain. This result was consistent with the activity of AFP and caspofungin [10], but differed to the function of PAF, where no CWIP activation and no induction of cell wall biosynthesis genes occurred [9].

Therefore, we conclude that AFP_{NN5353} triggers cell wall remodeling via Pkc/Mpk signalling. We further deduce from our data that similarities and differences exist in the molecular targets and the mode of action of antifungal proteins from filamentous fungi, e.g. AFP_{NN5353} and PAF - despite their homology. This phenomenon was also reported for other closely related antifungal proteins, such as the plant defensins MsDef1 and MtDef4 from *Medicago spp.* [44].

Apart from the activation of the CWIP, the perturbation of the Ca²⁺ homeostasis represents a major mechanistic function of antifungal proteins in sensitive fungi [17,18]. The intracellular Ca²⁺ response to AFP_{NN5353} in *A. niger* reflected that of the *Penicillium* antifungal protein PAF in *N. crassa* [17]. The rapid and sustained increase of the [Ca²⁺]_c resting level depended on a sustained influx of Ca²⁺ ions from the external medium. Moreover, the AFP_{NN5353} induced changes in the Ca²⁺ signature of mechanically perturbed *A. niger* cells further underlines the disruption of the Ca²⁺ response and homeostasis by AFP_{NN5353}. The addition of CaCl₂ to the growth medium reduced the susceptibility of *A. niger* towards the antifungal protein and decreased the AFP_{NN5353} specific rise in the [Ca²⁺]_c resting level. Both observations point towards an adaptive

response which is mediated most probably via Ca^{2+} signalling. First, high extracellular Ca^{2+} concentrations trigger chitin synthesis in *A. niger* and thereby confer increased protection against antifungal proteins as shown for AFP [15]. Second, it primes the Ca^{2+} homeostatic machinery to better maintain a low $[\text{Ca}^{2+}]_c$ resting level when challenged with the antifungal protein, e.g. by (i) the increase of the activity of existing Ca^{2+} pumps/transporters to counteract the $\text{AFP}_{\text{NN5353}}$ -specific intracellular Ca^{2+} perturbation, or (ii) the modulation of the expression of Ca^{2+} channels/pumps/exchangers [17]. The former hypothesis (i) might be supported by the observation that the addition of CaCl_2 only 10 min before *A. niger* was challenged with $\text{AFP}_{\text{NN5353}}$ restored the low $[\text{Ca}^{2+}]_c$ resting level. However, the perturbation of the Ca^{2+} homeostasis by a sustained elevation of the $[\text{Ca}^{2+}]_c$ resting level indicates that *A. niger* is not able to restore the low $[\text{Ca}^{2+}]_c$ resting level after exposure to $\text{AFP}_{\text{NN5353}}$ and this might trigger programmed cell death (PCD) on the long term as it was shown to occur in *A. nidulans* in response to the *P. chrysogenum* PAF [34].

Since AFP was shown to cause membrane permeabilization [21], the influx of Ca^{2+} might be due to changes in membrane permeability for this ion, if not the formation of pores. However, our staining experiments with CMFDA and PI exclude this possibility at least in the first 10 min of exposure to $\text{AFP}_{\text{NN5353}}$ when the $[\text{Ca}^{2+}]_c$ resting level reaches its maximum. This result is further corroborated by the fact that higher external concentrations of Ca^{2+} reduced the $\text{AFP}_{\text{NN5353}}$ specific rise in $[\text{Ca}^{2+}]_c$ resting level which - in our opinion - would not occur with leaky membranes. However, we do not exclude changes in membrane permeability at longer exposure times to this antifungal protein and more studies are needed to answer this question.

Finally, we observed that the internalization of $\text{AFP}_{\text{NN5353}}$ is characteristic for sensitive but not resistant moulds. A lack of binding of $\text{AFP}_{\text{NN5353}}$ to insensitive fungi might point towards the absence or inaccessibility of a putative interacting molecule at the cell surface. $\text{AFP}_{\text{NN5353}}$ localized to the cytoplasm of target fungi only when actin filaments were formed. This is in agreement with the endocytotic uptake and intracellular localization of the *P. chrysogenum* antifungal protein PAF in sensitive filamentous fungi [14,45]. Importantly, we observed that $\text{AFP}_{\text{NN5353}}$ was internalized by hyphae even under sub-inhibitory concentrations (0.2 $\mu\text{g}/\text{ml}$ for *A. nidulans*) which suggests that a threshold concentration is required to cause severe growth defects in target fungi.

The presence of high concentrations of extracellular Ca^{2+} counteracted $\text{AFP}_{\text{NN5353}}$ uptake. This finding parallels well with the report of [20] that the presence of cations, such as Ca^{2+} , interfered with the binding of AFP to the surface of *F. oxysporum* and with our observations made

with the *Penicillium* PAF (unpublished data). One possible explanation might be that extracellular Ca^{2+} ions compete with $\text{AFP}_{\text{NN5353}}$ for the same molecular target on the fungal surface which might represent a first binding receptor or even a "gate" for protein uptake [20,21] or, alternatively, that the interacting target is repressed under these conditions [17]. An additional explanation might be that the primary cell-surface localized $\text{AFP}_{\text{NN5353}}$ target might be masked due to a Ca^{2+} -dependent stimulation of chitin synthesis and cell wall remodeling as recently observed for AFP in *A. niger* [15]. This further suggests that the activation of the CWIP and the *agsA* induction does not mediate sufficient resistance to survive the toxic effects of $\text{AFP}_{\text{NN5353}}$. Instead, according to the "damage-response framework of AFP-fungal interactions" [15], the chitin response might represent the better strategy for fungi to survive the antifungal attack.


Conclusions

Based on the growth inhibitory activity, antifungal proteins like $\text{AFP}_{\text{NN5353}}$ can be well considered as promising candidates for future antimycotic drug developments. However, for biotechnological exploitation, the detailed knowledge on the mode of action is demanded. Our study shows that the detrimental effects caused by the *A. giganteus* antifungal protein $\text{AFP}_{\text{NN5353}}$ in sensitive target aspergilli are based on the interaction of this protein with more than one signalling pathway. In Figure 7, we present a tentative working model. The toxicity of $\text{AFP}_{\text{NN5353}}$ is mediated via PkcA/MpkA signalling which occurs independently from RhoA. Instead, so far unidentified RhoA-GAP effector molecules might contribute to $\text{AFP}_{\text{NN5353}}$ toxicity. The activation of the CWIP by $\text{AFP}_{\text{NN5353}}$ induces the *agsA* gene expression which is, however, insufficient to counteract toxicity of the protein. Furthermore, $\text{AFP}_{\text{NN5353}}$ leads to an immediate and significant increase of the $[\text{Ca}^{2+}]_c$ resting level in the cell. This sustained perturbation of the Ca^{2+} homeostasis could lead to PCD [17,34]. The presence of extracellular Ca^{2+} neutralizes the toxic effects of $\text{AFP}_{\text{NN5353}}$ and improves the resistance of the target organism possibly by decreasing the elevated $[\text{Ca}^{2+}]_c$ resting level and stimulating the fortification of the cell wall by the induction of *chsD* expression as shown for AFP [15]. Further investigations are in progress to clarify how these pathways are interconnected and interfere with each other on the molecular level.

Methods

Strains, Media and Chemicals

Fungal strains used in this study are listed in Table 5. All strains were obtained from the culture collections FGSC, ATCC, CBS, from the Institute of Microbiology, Division of Systematics, Taxonomy and Evolutionary



Biology at the Leopold Franzens University of Innsbruck, or the strain collection of the Department of Biotechnology, National Institute of Chemistry, Ljubljana, Slovenia. Unless otherwise stated, all fungi were grown in complete medium (CM) [19] with the respective supplements [28,38]. R153 and *alcA*-PkcA were grown in defined minimal medium (MM) according to [26]. Ca²⁺ response experiments were performed in Vogels medium [46]. For experiments with CaCl₂ supplementation, the KH₂PO₄ concentration of the culture media was reduced from 37 mM to 10 mM to avoid precipitation of supplemental Ca²⁺ and these media were called CM* and Vogels*. Chemicals were purchased from Sigma. AFP_{NN5353} and polyclonal rabbit anti-AFP_{NN5353} antibody were generous gifts from

Mogens T. Hansen, Novozymes, Denmark. The antifungal protein was isolated from *A. giganteus* strain A3274 (CBS 526.65), purified and analyzed by HPLC as described in the patent application WO94/01459 [47].

Growth inhibition assays

Antifungal activity assays were performed in 96-well plates in CM or Vogels medium inoculated with 1 × 10⁴ conidia/ml and supplemented with various concentrations of AFP_{NN5353} or with equivalent amounts of buffer (untreated controls). Fungal growth was monitored microscopically with an Olympus CK40 microscope equipped with a Zeiss MRc digital camera and the growth rates were determined spectrophotometrically as described previously [19]. Alternatively, 2 × 10³ conidia were spotted in

Table 5 Fungal strains used in this study.

Strain	Relevant genotype	Source or reference
<i>A. flavus</i> ATCC 9643	wild type	ATCC
<i>A. fumigatus</i> ATCC 46645	wild type	ATCC
<i>A. giganteus</i> AG 090701	wild type isolate	Institute of Microbiology
<i>A. nidulans</i>		
FGSC A4	Glasgow wild type (veA ⁺); velvet mutant	FGSC
R153	<i>wA2; pyroA4</i>	[26]
<i>alcA-PkcA</i>	<i>pyrG89::pyr4 alcA(p)::pkcAΔp</i>	[26]
GR5	<i>pyrG89; wA3; pyroA4</i>	[28]
RhoA ^{G14V}	GR5 + pGG2 (<i>rhoA</i> ^{G14V}) and pRG3AMA1 (co-transformation plasmid)	[28]
RhoA ^{E40I}	GR5 + pGG5 (<i>rhoA</i> ^{E40I}) and pRG3AMA1 (co-transformation plasmid)	[28]
<i>ΔmpkA</i>	<i>ΔmpkA</i>	[38]
<i>A. niger</i>		
CBS 120.49	wild type	CBS
A533	<i>cspA1, aeqS, amdS</i> ⁺ (pAEQS1-15)	[31]
RD6.47	<i>P agsA::h2b::egfp::Ttrpc</i>	[10]
<i>A. terreus</i> 304	wild type isolate	Institute of Microbiology
<i>Botrytis cinerea</i> BC 080801	wild type isolate	Institute of Microbiology
<i>Fusarium oxysporum</i> FO 240901	wild type isolate	Institute of Microbiology
<i>F. sambucinum</i> FS 210901	wild type isolate	Institute of Microbiology
<i>Gliocladium roseum</i> GR 210901	wild type isolate	Institute of Microbiology
<i>M. circinelloides</i> MC 080801	wild type isolate	Institute of Microbiology
<i>M. genevensis</i> MG 080801	wild type isolate	Institute of Microbiology
<i>P. chrysogenum</i> ATCC 10002	wild type	ATCC
<i>Trichoderma koningii</i> TC 060901	wild type isolate	Institute of Microbiology
<i>Neurospora crassa</i> FGSC 2489	wild type	FGSC

5 μl aliquots on appropriately supplemented agar plates. The plates were then incubated at 37°C for up to 72 h. Every 24 h, the plates were photographed and the colony diameters were determined. All assays were performed as technical triplicates and biological duplicates.

Analysis of the induction of the *agsA* expression by a GFP-based reporter system

The *A. niger* reporter strain RD6.47 carries the *agsA* promoter fused to a nucleus-targeted GFP (H2B::eGFP) [27]. Activation of the CWIP can be monitored by the increase in nuclear fluorescence. Analysis of the activation of the *agsA* promoter by 10-100 μg/ml AFP_{NN5353} was performed as described in [10]. As a positive control, caspofungin at a concentration of 10 μg/ml was used. Fluorescence images were taken from coverslips observed with an Axioplan 2 microscope (Zeiss) equipped with a Sony DKC-5000 digital camera.

Fluorescence staining

Indirect immunofluorescence staining

A. nidulans was grown over night on glass cover slips at 30°C in CM. They were further incubated for 90 min in

the presence or absence (controls) of 0.2 μg/ml AFP_{NN5353}. The samples were stained as described previously [14] and incubated with rabbit-anti-AFP_{NN5353} antibody (1:2.500, Novozymes, Denmark) for at least 60 min. Immunocomplexes were detected with FITC-conjugated swine-anti-rabbit IgG (1:40, DAKO, Germany). All samples were embedded in Vectashield mounting medium (Vector Laboratories, Burlingame, USA). Microscopy was done with a Zeiss Axioplan fluorescence microscope or a Zeiss confocal laser scanning microscope as described in [14].

For incubation with latrunculin B (Sigma, Austria), samples were treated with 0.2 μg/ml AFP_{NN5353} and 10 μg/ml latrunculin B for 80 min. As a control, samples were treated with DMSO to exclude artifacts evoked by the dissolvent of latrunculin B.

For detection of AFP_{NN5353} in the presence of elevated concentrations of CaCl₂, fungi were grown in CM* medium and then treated with 0.2 μg/ml AFP_{NN5353} in the presence of 10 mM CaCl₂ for 90 min.

Analysis of membrane permeabilization and cell viability

To determine if AFP_{NN5353} permeabilized the plasma membrane of *A. niger* germlings, we used a combination

of propidium iodide (PI) and fluorescein diacetate (cell tracker, CMFDA green, Invitrogen) according to [48]. Twelve h old *A. niger* germlings were grown in Vogels medium and pretreated with the two dyes (final conc. 5 µg/ml each) for 15 min before AFP_{NN5353} was added to a final concentration of 20 µg/ml. Samples without AFP_{NN5353} served as controls for positive CMFDA staining, while ethanol (70%) was used to permeabilize the membrane for positive PI staining.

Analysis of the calcium response to AFP_{NN5353} application
10⁵ conidia/ml of the *A. niger* strain A533 expressing codon optimized aequorin were grown in Vogels* medium containing 10 µM coelenterazine (Biosynth, Switzerland) at 30°C for twelve h in the dark. The [Ca²⁺]_c resting level and mechanical perturbation experiments and the calibration of [Ca²⁺]_c were performed as described in [17].

Additional material

Additional file 1: The expression of nucleus-targeted GFP under the control of the *agsA* promoter in *A. niger* in response to cell wall interfering substances. Differential interfering contrast images and corresponding fluorescence images of *A. niger* RD6.47 indicate the expression of a nucleus-targeted GFP under the control of the *A. niger* *agsA* promoter. Five h old germlings were (A) left untreated (negative control), (B) treated with 50 µg/ml AFP_{NN5353} and (C) with 10 µg/ml caspofungin (positive control) as described in Materials and Methods. Scale bar, 20 µm.

Additional file 2: Viability staining of *A. niger* germlings after AFP_{NN5353} exposure. Twelve h old *A. niger* germlings were stained with fluorescein diacetate (CMFDA, middle panels) and propidium iodide (right panels). The left panels show the respective light micrographs. All samples were pretreated with the dyes for 15 min before 20 µg/ml AFP_{NN5353} was added (B). Controls remained untreated (A) or were exposed to 70% ethanol (C). Scale bar, 50 µm.

Acknowledgements

We thank Mogens T. Hansen (Novozymes, Denmark) for the generous gift of AFP_{NN5353} and the polyclonal rabbit anti-AFP_{NN5353} antibody. We gratefully acknowledge Renate Weiler-Görz for technical assistance. This study was financially supported by the Austrian Science Fund FWF (P19970-B11) and the Österreichischer Austauschdienst ÖAD (Wissenschaftlich-Technische Zusammenarbeit Österreich und Slowenien, S115/2009).

Author details

¹Biocenter, Division of Molecular Biology, Innsbruck Medical University, Fritz-Pregl Strasse 3, Innsbruck, A-6020, Austria. ²Department of Biotechnology, National Institute of Chemistry, Hajdrihova 19, Ljubljana, SI-1000, Slovenia. ³Excellent NMR, Future Innovation for Sustainable Technologies Centre of Excellence, Hajdrihova 19, Ljubljana, SI-1000, Slovenia. ⁴Department of Applied and Molecular Microbiology, Institute of Biotechnology, Berlin University of Technology, Gustav-Meyer-Allee 25, Berlin, D-13355, Germany. ⁵Department of Hygiene, Microbiology and Social Medicine, Innsbruck Medical University, Fritz-Pregl Strasse 3, Innsbruck, A-6020, Austria.

Authors' contributions

UB carried out the growth inhibition assays, the indirect immunofluorescence stainings, the Ca²⁺ measurements and the calculations to convert the luminescence units into the [Ca²⁺]_c levels. She also performed the statistical analysis and helped to draft the manuscript. MB

contributed the *A. niger* A533 strain, helped with the Ca²⁺ measurements and participated in the design of the study. AE contributed to the indirect immunofluorescence stainings. VM contributed the *A. niger* RD6.47 strain and performed the *agsA* induction assays. FM conceived of the study, participated in its design and coordination and drafted the manuscript. All authors read and approved the final manuscript.

Received: 25 May 2011 Accepted: 23 September 2011

Published: 23 September 2011

References

1. Hancock RE, Scott MG: The role of antimicrobial peptides in animal defenses. *Proc Natl Acad Sci USA* 2000, **97**(16):8856-8861.
2. Kamysz W, Okroj M, Lukasiak J: Novel properties of antimicrobial peptides. *Acta Biochim Pol* 2003, **50**(2):461-469.
3. Aerts AM, Francois IE, Cammue BP, Thevissen K: The mode of antifungal action of plant, insect and human defensins. *Cell Mol Life Sci* 2008, **65**(13):2069-2079.
4. Gupte MD, Kulkarni PR: A study of antifungal antibiotic production by *Streptomyces chattanoogensis* MTCC 3423 using full factorial design. *Letts Appl Microbiol* 2002, **35**(1):22-26.
5. Geisen R: *P. nalgiovense* carries a gene which is homologous to the *paf* gene of *P. chrysogenum* which codes for an antifungal peptide. *Int J Food Microbiol* 2000, **62**(1-2):95-101.
6. Lee GD, Shin SY, Maeng CY, Jin ZZ, Kim KL, Hahn KS: Isolation and characterization of a novel antifungal peptide from *Aspergillus niger*. *Biochem Biophys Res Commun* 1999, **263**:646-651.
7. Marx F, Haas H, Reindl M, Stoffer G, Lottspeich F, Redl B: Cloning, structural organization and regulation of expression of the *Penicillium chrysogenum paf* gene encoding an abundantly secreted protein with antifungal activity. *Gene* 1995, **167**(1-2):167-171.
8. Wnendt S, Ulbrich N, Stahl U: Molecular cloning, sequence analysis and expression of the gene encoding an antifungal-protein from *Aspergillus giganteus*. *Curr Genet* 1994, **25**(6):519-523.
9. Binder U, Oberparleiter C, Meyer V, Marx F: The antifungal protein PAF interferes with PKC/MPK and cAMP/PKA signalling of *Aspergillus nidulans*. *Mol Microbiol* 2010, **75**(2):294-307.
10. Hagen S, Marx F, Ram AF, Meyer V: The antifungal protein AFP from *Aspergillus giganteus* inhibits chitin synthesis in sensitive fungi. *Appl Environ Microbiol* 2007, **73**(7):2128-2134.
11. Lobo DS, Pereira IB, Fragel-Madeira L, Medeiros LN, Cabral LM, Faria J, Bellio M, Campos RC, Linden R, Kurtenbach E: Antifungal *Pisum sativum* defensin 1 interacts with *Neurospora crassa* cyclin F related to the cell cycle. *Biochemistry* 2007, **46**(4):987-996.
12. Marx F, Binder U, Leiter E, Pocs I: The *Penicillium chrysogenum* antifungal protein PAF, a promising tool for the development of new antifungal therapies and fungal cell biology studies. *Cell Mol Life Sci* 2008, **65**(3):445-454.
13. Moreno AB, Penas G, Rufat M, Bravo JM, Estopa M, Messeguer J, San Segundo B: Pathogen-induced production of the antifungal AFP protein from *Aspergillus giganteus* confers resistance to the blast fungus *Magnaporthe oryzae* in transgenic rice. *Mol Plant Microbe Interact* 2005, **18**(9):960-972.
14. Oberparleiter C, Kaiserer L, Haas H, Ladurner P, Andratsch M, Marx F: Active internalization of the *Penicillium chrysogenum* antifungal protein PAF in sensitive aspergilli. *Antimicrob Agents Chemother* 2003, **47**(11):3598-3601.
15. Ouedraogo JP, Hagen S, Spielvogel A, Engelhardt S, Meyer V: Survival strategies of yeast and filamentous fungi against the antifungal protein AFP. *J Biol Chem* 2011, **286**(16):13859-13868.
16. Fujioka T, Mizutani O, Furukawa K, Sato N, Yoshimi A, Yamagata Y, Nakajima T, Abe K: MpkA-Dependent and -independent cell wall integrity signaling in *Aspergillus nidulans*. *Eukaryot Cell* 2007, **6**(8):1497-1510.
17. Binder U, Chu M, Read ND, Marx F: The antifungal activity of the *Penicillium chrysogenum* protein PAF disrupts calcium homeostasis in *Neurospora crassa*. *Eukaryot Cell* 2010, **9**(9):1374-1382.
18. Thevissen K, Ghazi A, De Samblanx GW, Brownlee C, Osborn RW, Broekaert WF: Fungal membrane responses induced by plant defensins and thionins. *J Biol Chem* 1996, **271**(25):15018-15025.
19. Kaiserer L, Oberparleiter C, Weiler-Görz R, Burgstaller W, Leiter E, Marx F: Characterization of the *Penicillium chrysogenum* antifungal protein PAF. *Arch Microbiol* 2003, **180**(3):204-210.

20. Martin-Urdiroz M, Martinez-Rocha AL, Di Pietro A, Martinez-del-Pozo A, Roncero MI: **Differential toxicity of antifungal protein AFP against mutants of *Fusarium oxysporum***. *Int Microbiol* 2009, **12**(2):115-121.
21. Theis T, Wedde M, Meyer V, Stahl U: **The antifungal protein from *Aspergillus giganteus* causes membrane permeabilization**. *Antimicrob Agents Chemother* 2003, **47**(2):588-593.
22. Wnendt S, Felske-Zech H, Henze PP, Ulbrich N, Stahl U: **Characterization of the gene encoding alpha-sarcin, a ribosome-inactivating protein secreted by *Aspergillus giganteus***. *Gene* 1993, **124**(2):239-244.
23. Meyer V: **A small protein that fights fungi: AFP as a new promising antifungal agent of biotechnological value**. *Appl Microbiol Biotechnol* 2008, **78**(1):17-28.
24. Levin DE: **Cell wall integrity signaling in *Saccharomyces cerevisiae***. *Microbiol Mol Biol Rev* 2005, **69**(2):262-291.
25. Damveld RA, Arentshorst M, Franken A, vanKuyk PA, Klis FM, van den Hondel CA, Ram AF: **The *Aspergillus niger* MADS-box transcription factor RlmA is required for cell wall reinforcement in response to cell wall stress**. *Mol Microbiol* 2005, **58**(1):305-319.
26. Ronen R, Sharon H, Levdansky E, Romano J, Shadkchan Y, Oshero N: **The *Aspergillus nidulans* *pkcA* gene is involved in polarized growth, morphogenesis and maintenance of cell wall integrity**. *Curr Genet* 2007, **51**(5):321-329.
27. Meyer V, Damveld RA, Arentshorst M, Stahl U, van den Hondel CA, Ram AF: **Survival in the presence of antifungals: genome-wide expression profiling of *Aspergillus niger* in response to sublethal concentrations of caspofungin and fenpropimorph**. *J Biol Chem* 2007, **282**(45):32935-32948.
28. Guest GM, Lin X, Momany M: ***Aspergillus nidulans* RhoA is involved in polar growth, branching, and cell wall synthesis**. *Fungal Genet Biol* 2004, **41**(1):13-22.
29. Terras FR, Schoofs HM, De Bolle MF, Van Leuven F, Rees SB, Vanderleyden J, Cammue BP, Broekaert WF: **Analysis of two novel classes of plant antifungal proteins from radish (*Raphanus sativus* L.) seeds**. *J Biol Chem* 1992, **267**(22):15301-15309.
30. Terras FR, Torrekens S, Van Leuven F, Osborn RW, Vanderleyden J, Cammue BP, Broekaert WF: **A new family of basic cysteine-rich plant antifungal proteins from *Brassicaceae* species**. *FEBS Lett* 1993, **316**(3):233-240.
31. Bencina M, Legisa M, Read ND: **Cross-talk between cAMP and calcium signalling in *Aspergillus niger***. *Mol Microbiol* 2005, **56**(1):268-281.
32. Nelson G, Kozlova-Zwinderman O, Collis AJ, Knight MR, Fincham JR, Stanger CP, Renwick A, Hessing JG, Punt PJ, van den Hondel CA, Read ND: **Calcium measurement in living filamentous fungi expressing codon-optimized aequorin**. *Mol Microbiol* 2004, **52**(5):1437-1450.
33. Moreno AB, Martinez Del Pozo A, San Segundo B: **Biotechnologically relevant enzymes and proteins. Antifungal mechanism of the *Aspergillus giganteus* AFP against the rice blast fungus *Magnaporthe grisea***. *Appl Microbiol Biotechnol* 2006, **72**(5):883-895.
34. Leiter E, Szappanos H, Oberparleiter C, Kaiserer L, Csernoch L, Pusztahelyi T, Emri T, Pócsi I, Salvenmoser W, Marx F: **Antifungal protein PAF severely affects the integrity of the plasma membrane of *Aspergillus nidulans* and induces an apoptosis-like phenotype**. *Antimicrob Agents Chemother* 2005, **49**(6):2445-2453.
35. Morton WM, Ayscough KR, McLaughlin PJ: **Latrunculin alters the actin-monomer subunit interface to prevent polymerization**. *Nat Cell Biol* 2000, **2**(6):376-378.
36. Conner SD, Schmid SL: **Regulated portals of entry into the cell**. *Nature* 2003, **422**(6927):37-44.
37. Lamaze C, Fujimoto LM, Yin HL, Schmid SL: **The actin cytoskeleton is required for receptor-mediated endocytosis in mammalian cells**. *J Biol Chem* 1997, **272**(33):20332-20335.
38. Bussink HJ, Osmani SA: **A mitogen-activated protein kinase (MPKA) is involved in polarized growth in the filamentous fungus, *Aspergillus nidulans***. *FEMS Microbiol Lett* 1999, **173**(1):117-125.
39. Ichinomiya M, Uchida H, Koshi Y, Ohta A, Horiuchi H: **A protein kinase C-encoding gene, *pkcA*, is essential to the viability of the filamentous fungus *Aspergillus nidulans***. *Biosci Biotechnol Biochem* 2007, **71**(11):2787-2799.
40. Reinoso-Martin C, Schuller C, Schuetzer-Muehlbauer M, Kuchler K: **The yeast protein kinase C cell integrity pathway mediates tolerance to the antifungal drug caspofungin through activation of Sit2p mitogen-activated protein kinase signaling**. *Eukaryot Cell* 2003, **2**(6):1200-1210.
41. Igual JC, Johnson AL, Johnston LH: **Coordinated regulation of gene expression by the cell cycle transcription factor Swi4 and the protein kinase C MAP kinase pathway for yeast cell integrity**. *EMBO J* 1996, **15**(18):5001-5013.
42. Jung US, Levin DE: **Genome-wide analysis of gene expression regulated by the yeast cell wall integrity signalling pathway**. *Mol Microbiol* 1999, **34**(5):1049-1057.
43. Kuranda K, Leberre V, Sokol S, Palamarczyk G, Francois J: **Investigating the caffeine effects in the yeast *Saccharomyces cerevisiae* brings new insights into the connection between TOR, PKC and Ras/cAMP signalling pathways**. *Mol Microbiol* 2006, **61**(5):1147-1166.
44. Ramamoorthy V, Zhao X, Snyder AK, Xu JR, Shah DM: **Two mitogen-activated protein kinase signalling cascades mediate basal resistance to antifungal plant defensins in *Fusarium graminearum***. *Cell Microbiol* 2007, **9**(6):1491-1506.
45. Batta G, Barna T, Gaspari Z, Sandor S, Kover KE, Binder U, Sarg B, Kaiserer L, Chhillar AK, Eigentler A, Leiter É, Hgedüs N, Pócsi I, Lindner H, Marx F: **Functional aspects of the solution structure and dynamics of PAF - a highly-stable antifungal protein from *Penicillium chrysogenum***. *FEBS J* 2009, **276**(10):2875-2890.
46. Vogel HJA: **Convenient growth medium for *Neurospora* (Medium N)**. *Microbiological Genetics Bulletin* 1956, **13**:42-43.
47. Pedersen AH, Halkier T, Nielsen BA, Lange L, Mikkelsen JM, Rasmussen G, Hansen MT: **A fungicidally active compound Novo Nordisk A/S, Denmark; Patent WO 94/01459; 1994**.
48. Palma-Guerrero J, Huang IC, Jansson HB, Salinas J, Lopez-Llorca LV, Read ND: **Chitosan permeabilizes the plasma membrane and kills cells of *Neurospora crassa* in an energy dependent manner**. *Fungal Genet Biol* 2009, **46**(8):585-594.

doi:10.1186/1471-2180-11-209

Cite this article as: Binder et al.: The *Aspergillus giganteus* antifungal protein AFP_{NNS353} activates the cell wall integrity pathway and perturbs calcium homeostasis. *BMC Microbiology* 2011 **11**:209.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

