

RESEARCH ARTICLE

Open Access

Alleles of the homologous recombination gene, *RAD59*, identify multiple responses to disrupted DNA replication in *Saccharomyces cerevisiae*

Lauren C Liddell^{1,2}, Glenn M Manthey¹, Shannon N Owens³, Becky XH Fu⁴ and Adam M Bailis^{1,2*}

Abstract

Background: In *Saccharomyces cerevisiae*, Rad59 is required for multiple homologous recombination mechanisms and viability in DNA replication-defective *rad27* mutant cells. Recently, four *rad59* missense alleles were found to have distinct effects on homologous recombination that are consistent with separation-of-function mutations. The *rad59-K166A* allele alters an amino acid in a conserved α -helical domain, and, like the *rad59* null allele diminishes association of Rad52 with double-strand breaks. The *rad59-K174A* and *rad59-F180A* alleles alter amino acids in the same domain and have genetically similar effects on homologous recombination. The *rad59-Y92A* allele alters a conserved amino acid in a separate domain, has genetically distinct effects on homologous recombination, and does not diminish association of Rad52 with double-strand breaks.

Results: In this study, *rad59* mutant strains were crossed with a *rad27* null mutant to examine the effects of the *rad59* alleles on the link between viability, growth and the stimulation of homologous recombination in replication-defective cells. Like the *rad59* null allele, *rad59-K166A* was synthetically lethal in combination with *rad27*. The *rad59-K174A* and *rad59-F180A* alleles were not synthetically lethal in combination with *rad27*, had effects on growth that coincided with decreased ectopic gene conversion, but did not affect mutation, unequal sister-chromatid recombination, or loss of heterozygosity. The *rad59-Y92A* allele was not synthetically lethal when combined with *rad27*, stimulated ectopic gene conversion and heteroallelic recombination independently from *rad27*, and was mutually epistatic with *srs2*. Unlike *rad27*, the stimulatory effect of *rad59-Y92A* on homologous recombination was not accompanied by effects on growth rate, cell cycle distribution, mutation, unequal sister-chromatid recombination, or loss of heterozygosity.

Conclusions: The synthetic lethality conferred by *rad59* null and *rad59-K166A* alleles correlates with their inhibitory effect on association of Rad52 with double-strand breaks, suggesting that this may be essential for rescuing replication lesions in *rad27* mutant cells. The *rad59-K174A* and *rad59-F180A* alleles may fractionally reduce this same function, which proportionally reduced repair of replication lesions by homologous recombination and growth rate. In contrast, *rad59-Y92A* stimulates homologous recombination, perhaps by affecting association of replication lesions with the Rad51 recombinase. This suggests that Rad59 influences the rescue of replication lesions by multiple recombination factors.

Keywords: Homologous recombination, *Saccharomyces cerevisiae*, DNA replication, Genome stability, Loss of heterozygosity

* Correspondence: abailis@coh.org

¹Department of Molecular and Cellular Biology, Beckman Research Institute of the City of Hope, 91010 Duarte, CA, USA

²Irell & Manella Graduate School of Biological Sciences, Beckman Research Institute of the City of Hope, 91010 Duarte, CA, USA

Full list of author information is available at the end of the article

Background

In *Saccharomyces cerevisiae*, defective DNA replication stimulates homologous recombination (HR), suggesting that the lesions that accumulate following replication failure are substrates for HR [1-11]. Rad27 is a structure-specific endonuclease [12] required for completion of lagging strand synthesis [13], and has also been implicated in base excision repair [14], and double-strand break repair by non-homologous end joining [15]. Loss of Rad27 leads to accumulation of single-stranded gaps or nicks on daughter DNA strands [2,16]. Collision of replication forks with these lesions results in fork collapse and generation of double-strand breaks (DSB) [8,17] that can stimulate HR. Importantly, concomitant loss of Rad27 and components of the HR apparatus leads to synthetic lethality [18-20]. These observations implicate HR in repair of DSBs that accumulate in the absence of Rad27. Failure to repair DSBs leads to chromosome loss [21] that is greatly stimulated in *rad27* null mutant cells [8], suggesting that the essential role for the HR apparatus in *rad27* mutants may be prevention of lethal levels of chromosome loss.

RAD59 encodes a protein that augments the ability of Rad52, the central HR protein in yeast [22,23], to anneal complementary DNA strands *in vitro* [24], and both are required for viability in *rad27* null mutant cells [19,20]. *RAD59* and *RAD52* are also required to repair DSBs by single-strand annealing (SSA) [21,25-28], and HR between inverted repeats by an annealing-dependent template switch at stalled replication forks [29-31]. Since *RAD59* exerts much of its effect on HR with *RAD52* [21,32,33], the function of *RAD59* required in the absence of *RAD27* may be in collaboration with *RAD52*.

The purpose of the current study was to explore the function of *RAD59* required for the viability of *rad27* null mutant cells. We investigated how four *rad59* mutations previously characterized with respect to their effects on SSA [21,27], affected survivorship when combined with a *rad27* null mutation. We found that *rad59-K166A*, which alters an amino acid in a conserved, putative α -helical domain [27,34,35], was synthetically lethal in combination with *rad27*. Because *rad59-K166A* diminishes association of Rad52 with DSBs [21], this may be a function required for the viability of *rad27* null mutant cells. The *rad59-K174A* and *rad59-F180A* mutations, which alter amino acids in the same α -helical domain, and have genetically similar effects on SSA [21], were not synthetically lethal with *rad27*, but resulted in distinct effects on growth that correlated with their degree of inhibition of HR. This strongly implicates *RAD59*-dependent HR as a requirement for viability in *rad27* null mutant cells. The *rad59-Y92A* mutation, which alters an amino acid in a separate, conserved loop domain and confers genetically distinct effects on SSA [27,34] was not synthetically lethal with

rad27, and had a stimulatory effect on HR. This effect was genetically equivalent to that of a null allele of *SRS2*, which encodes a helicase that disassembles Rad51-DNA filaments [36,37], suggesting that Rad59 may affect association of Rad51 with replication lesions. The distinct effects of the *rad59* alleles suggest that Rad59 possesses multiple, discrete roles in responding to the consequences of dysfunctional replication.


Results

The *rad59* mutant alleles display distinct effects on survival and growth in cells defective for lagging strand synthesis

To further explore the function of *RAD59* required for viability in *rad27* null mutant cells, the effects of combining the *rad27::LEU2* allele with the various *rad59* alleles were determined by examining their ability to yield viable spores upon co-segregation in genetic crosses. The various *RAD27/rad27::LEU2 RAD59/rad59* double heterozygotes were sporulated and tetrads dissected onto rich medium (Figure 1). As observed previously, the *rad27::LEU2* and *rad59::LEU2* alleles did not appear together in any of the colonies arising from the spores, consistent with synthetic lethality [19,20]. The *rad59-K166A* allele, which alters a conserved lysine in the region of Rad59 that corresponds to the α -helical domain of the $\beta - \beta - \beta - \alpha$ motif of human Rad52 (Additional file 1: Figure S1) [27,34,35] displayed the same failure to appear with the *rad27::LEU2* allele, indicative of synthetic lethality.

The *rad59-K174A* and *rad59-F180A* alleles alter conserved amino acids in the same putative α -helical domain as *rad59-K166A* but were able to form viable spores upon segregation with *rad27::LEU2* (Figure 1). Doubling time of the *rad27::LEU2 rad59-F180A* double mutant was a statistically significant ($p = 0.045$) 24% longer than that observed for the *rad27* single mutant, which correlated with a ratio of G1 to S + G2/M cells that was a statistically significant ($p = 0.0031$) 2.6-fold lower (Figure 2; Additional file 1: Table S2). In contrast, doubling time of the *rad27::LEU2 rad59-K174A* double mutant was not significantly different from that of the *rad27::LEU2* single mutant ($p = 0.71$) (Table 1; Additional file 1: Table S2).

The *rad59-Y92A* allele alters a conserved amino acid in another region of extensive conservation with Rad52 (Additional file 1: Figure S1) [27,34], and was observed to yield viable spores upon segregation with *rad27::LEU2* (Figure 1). While the colonies derived from the *rad27::LEU2 rad59-Y92A* double mutant spores sometimes appeared smaller than the *rad27::LEU2* single mutant colonies on dissection plates, neither the doubling times ($p = 0.707$) (Table 1; Additional file 1: Table S2), nor the ratios of G1 to S + G2/M cells ($p = 0.60$) (Figure 2, Additional file 1: Table S2) were significantly different for the *rad27::LEU2* single and *rad27::LEU2 rad59-Y92A*



double mutant strains. This suggests that germination of *rad27::LEU2 rad59-Y92A* double mutant spores may sometimes take longer than *rad27::LEU2* single mutant spores. We did not observe significant effects of the tested *rad59* missense alleles on doubling time ($p > 0.15$) (Table 1; Additional file 1: Table S2), or cell cycle distribution ($p > 0.50$) (Figure 2; Additional file 1: Table S2) in cells that possessed a wild-type *RAD27* gene. Since all four *rad59* missense mutations support steady-state levels of Rad59 that are comparable to wild-type [27], their effects on viability and growth when combined with *rad27::LEU2* cannot be attributed to changes in the level of Rad59 in the cell. Altogether, these observations suggest that *RAD59* plays a critical role in determining the growth characteristics of cells defective for lagging strand synthesis.

The *rad59* alleles affect a *RAD51*-dependent mechanism for repairing replication lesions

The central strand exchange factor, Rad51 [38], is often required for mechanisms of HR that require *RAD59*, including those involved in spontaneous HR between inverted and unlinked repeat sequences [39,40]. Like *RAD59*, an intact *RAD51* gene is necessary for viability in *rad27::LEU2* mutant cells [18-20], suggesting that *RAD51*-dependent HR plays a critical role in responding to replication lesions. Accordingly, loss of *RAD27* results in increases in HR events that require *RAD51* [18]. We used an assay that measures spontaneous ectopic gene conversion involving unlinked, mutant alleles of the

SAM1 gene [41] to examine effects of the *rad27::LEU2* mutation on HR in haploid strains (Figure 3A). Loss of *RAD27* resulted in a dramatic, 4,700-fold increased rate of ectopic gene conversion (Figure 3B; Additional file 1: Table S2), indicating that accumulation of replication lesions can greatly stimulate HR between unlinked sequences.

The robust stimulatory effect of the loss of the *RAD27* gene on ectopic gene conversion suggested that it could be used for examining the relationship between HR, and growth in the viable *rad27 rad59* double mutants. As observed previously [40], the *rad59::LEU2* mutation conferred a statistically significant 2.7-fold reduction in the rate of ectopic gene conversion (Figure 3B; Additional file 1: Table S2), confirming that *RAD59* plays a role in spontaneous HR between unlinked repeats. While neither the *rad59-K174A* nor *rad59-F180A* mutations had significant effects on their own, they led to significant, 3.1- and 9.3-fold reductions in the stimulatory effect of the *rad27::LEU2* allele in the *rad27::LEU2 rad59-K174A* and *rad27::LEU2 rad59-F180A* double mutants (Figure 3C; Additional file 1: Table S2), suggesting that they confer defects in the utilization of replication lesions by HR.

In contrast to the *rad59-K174A* and *rad59-F180A* mutations, the *rad59-Y92A* mutation caused an 86-fold increased rate of spontaneous ectopic gene conversion (Figure 3B; Additional file 1: Table S2), and, when combined with the *rad27::LEU2* mutation, stimulated the rate of ectopic gene conversion by a statistically significant 7.7-fold over that observed in the *rad27::LEU2*




Table 1 Doubling times in wild-type and mutant haploid cells


Genotype	Doubling time (min)	95% confidence interval
Wild-type	111	99, 120
<i>rad59-Y92A</i>	119	97, 124
<i>rad59-K174A</i>	131	111, 147
<i>rad59-F180A</i>	112	99, 128
<i>rad27::LEU2</i>	164	137, 180
<i>rad27::LEU2 rad59-Y92A</i>	176	136, 195
<i>rad27::LEU2 rad59-K174A</i>	153	126, 177
<i>rad27::LEU2 rad59-F180A</i>	205	183, 230

Doubling times of freshly dissected segregants were determined as described in the Methods. Displayed for each genotype is the median doubling time and 95% confidence interval, determined from at least ten independent cultures.

single mutant (Figure 3B and C; Additional file 1: Table S2). The synergistically increased rate of ectopic gene conversion in the *rad27::LEU2 rad59-Y92A* double mutant is consistent with *rad59-Y92A* stimulating HR by a mechanism distinct from the accumulation of replication lesions that results from loss of *RAD27*.

The hyper-rec effects of the *rad59-Y92A* and *srs2::TRP1* alleles are genetically equivalent

Previous work indicating that *rad59-Y92A* decreases spontaneous *RAD51*-independent HR between directly repeated sequences [27] suggests that the stimulation of ectopic gene conversion is not due to accumulation of recombinogenic lesions. Ectopic gene conversion requires Rad51 to work after lesion formation to catalyze the strand invasion that begins the interaction between unlinked sequences that will repair the lesion [40,42]. If



stimulation of HR by *rad59-Y92A* is the result of changes subsequent to Rad51-DNA filament formation, loss of *RAD51* should abolish the stimulatory effect. The rate of ectopic gene conversion in the *rad51::LEU2 rad59-Y92A* double mutant was reduced 50-fold from wild-type, which was nearly identical to the rate in *rad51::LEU2* single mutant cells (Figure 3D; Additional

file 1: Table S2). Therefore, stimulation by *rad59-Y92A* requires formation of Rad51-DNA filaments.

Like the *rad59-Y92A* mutation, a null allele of the *SRS2* gene, which encodes a DNA helicase [43] that facilitates the disassembly of Rad51-DNA filaments [36,37], has been shown to stimulate spontaneous gene conversion between non-allelic sequences [44,45]. Consistent with

this, we observed a 31-fold increased rate of spontaneous ectopic gene conversion in an *srs2::TRP1* mutant (Figure 3D; Additional file 1: Table S2). As the effects of *srs2::TRP1* and *rad59-Y92A* were similar we examined ectopic gene conversion in the *srs2 rad59-Y92A* double mutant and observed a 38-fold increase over wild-type that was not significantly different from the rates in the *srs2::TRP1* or *rad59-Y92A* single mutants (Figure 3B and 3D; Additional file 1: Table S2). This indicates that *rad59-Y92A* and *srs2::TRP1* are mutually epistatic.

***rad59-Y92A* is a dominant mutation that stimulates *RAD51*-dependent HR between homologs**

RAD51-dependent HR between sister-chromatids and homologous chromosomes are thought to be the primary mechanisms for rescuing lesions, and supporting viability in *rad27* mutant cells [18-20]. One form of inter-homolog HR that requires *RAD51* is recombination between mutant alleles at the same locus, referred to as heteroallelic recombination [46]. Accordingly, the rate of spontaneous recombination between heteroalleles of the *SAM2* gene was reduced 10.5-fold in the *rad51::*

LEU2/rad51::LEU2 homozygote (Figure 4B; Additional file 1: Table S2). Consistent with its effect on ectopic gene conversion, loss of *RAD27* increased the rate of heteroallelic recombination 2,400-fold, confirming that accumulation of replication lesions robustly stimulates heteroallelic recombination [18].

Similar to its effect on ectopic gene conversion, we observed that *rad59-Y92A* increased the rate of heteroallelic recombination by 19-fold (Figure 4B; Additional file 1: Table S2). Interestingly, the effect of *rad59-Y92A* was dominant with respect to *RAD59*, as the rate in the *RAD59/rad59-Y92A* heterozygote was not significantly different from that in the *rad59-Y92A/rad59-Y92A* homozygote. Like with ectopic gene conversion, combining the *rad27::LEU2* and *rad59-Y92A* alleles in the *rad27/rad27 rad59-Y92A/rad59-Y92A* double homozygote had a synergistic effect on heteroallelic recombination, increasing the rate 25-fold over that observed in the *rad27::LEU2/rad27::LEU2* homozygote. This astonishing, 59,000-fold increased rate of heteroallelic recombination corresponds to a median frequency of recombination where 85% of the surviving colonies are recombinants.




Figure 4 The *rad59-Y92A* mutation has a dominant, hyper-rec effect on hetero-allelic recombination in diploid strains. **(A)** The spontaneous hetero-allelic recombination system: Diploid strains possessing a *sam2-ΔEcoRV* V-HOCs allele at the *SAM2* locus on one copy of chromosome IV, a *sam2-ΔSal I* allele on the other copy, and a *sam1::LEU2* allele at the *SAM1* locus on both copies of chromosome XII (not pictured) were grown to saturation in YPD medium supplemented with AdoMet, and plated onto medium lacking AdoMet to select for cells in which a recombination event generates a functional *SAM2* gene and an AdoMet prototrophic cell. Either reciprocal or non-reciprocal recombination events between *sam2-ΔEcoRV* V-HOCs and *sam2-ΔSal I* can generate AdoMet⁺ recombinants. The *sam1::LEU2* allele is missing sufficient information to recombine with the *sam2* alleles. The white bars indicate the positions of the *sam2* mutations. **(B)** Rates of heteroallelic recombination in wild-type, heterozygous and homozygous mutant strains. Rates were determined from a minimum of 10 independent cultures as described in the Methods. Fold decreases (-) and increases (+) from wild-type indicated in boxes.

The *rad59* alleles do not affect a variety of genome destabilizing processes stimulated by the accumulation of replication lesions

Loss of *RAD27* stimulates a variety of mutagenic and clastogenic events [8,16,18,47,48]. The rate of spontaneous mutation at the *CAN1* locus is greatly increased in *rad27::LEU2* mutant cells [8,18,49]. Characterization of these mutations revealed that the majority are short duplications flanked by short, directly repeated sequences that may be created by multiple HR mechanisms [18]. Our data confirm the previous analyses as we observed a 50-fold increased rate of spontaneous mutation at the *CAN1* locus in a *rad27::LEU2* mutant (Table 2; Additional file 1: Table S2). In contrast, the *rad59::LEU2*, *rad59-Y92A*, *rad59-K174A*, and *rad59-F180A* alleles did not have significant effects on the rate of *CAN1* mutation, nor did the missense alleles have significant effects when combined with the *rad27::LEU2* allele.

Loss of *RAD27* has been previously observed to strongly stimulate unequal sister chromatid recombination (USCR) (Additional file 1: Figure S2) [8,50]. We observed a 47-fold increased rate of USCR in *rad27::LEU2* cells (Table 2; Additional file 1: Table S2), confirming the previous results, while loss of *RAD51* had no significant effect. The *rad59::LEU2*, *rad59-Y92A*, *rad59-K174A*, and *rad59-F180A* alleles did not have significant effects on the rate of USCR, nor did the missense mutations have effects in combination with *rad27::LEU2*, suggesting that *RAD59* does not influence this mechanism of genome rearrangement.

Disrupting lagging strand synthesis by imposing a defect in the processivity of Pol δ , or loss of *RAD27*, was shown previously to substantially increase rates of loss of heterozygosity (LOH) by chromosome loss, and HR between homologs [2,8,10,11,18]. In the present analysis,

LOH was examined in diploid strains by simultaneously monitoring changes in the genetic state at three loci on chromosome V (*HXT13*, *CAN1* and *HOM3*) in order to separately determine rates of chromosome loss (reduction to hemizygoty at all three loci), terminal LOH (homozygosity at *HXT13* and *CAN1*), and interstitial LOH (homozygosity at *CAN1*) (Additional file 1: Figure S3; Table 3; Additional file 1: Table S2). Rates of all three events increased substantially in *rad27::LEU2/rad27::LEU2* homozygotes; chromosome loss increased 12-fold, terminal LOH increased 37-fold, and interstitial LOH increased 11-fold, strongly suggesting that replication lesions stimulate HR between homologs when they are repaired, and chromosome loss when they are not. Interestingly, we observed an 18-fold increase in the rate of chromosome loss in *rad51::LEU2/rad51::LEU2* homozygotes, consistent with a requirement for *RAD51* in the rescue of broken chromosomes. In contrast, loss of *RAD51* did not have significant effects on interstitial LOH or terminal LOH, indicating that these inter-chromosomal HR events do not require Rad51.

As observed above for mutation and USCR (Table 2; Additional file 1: Table S2), the *rad59-Y92A*, *rad59-K174A*, and *rad59-F180A* alleles had no significant effect on the rates of interstitial LOH, terminal LOH, and chromosome loss in the *rad59/rad59* single mutants, or in the double mutant combinations with the *rad27::LEU2* allele (Table 3; Additional file 1: Table S2). Similarly, *rad59::LEU2* had no significant effect on the rates of interstitial LOH and terminal LOH, but conferred a small (two-fold), statistically significant increase in chromosome loss. These data suggest that *RAD59* has little influence on these mechanisms of LOH.

Discussion

We have explored the role of *RAD59* in mediating responses to DNA lesions that accumulate in *rad27::LEU2* mutant cells, and found that it supports multiple, genetically separable functions. Like the *rad59::LEU2* allele, we found that the *rad59-K166A* allele, which alters a lysine in a conserved, putative α -helical domain (Additional file 1: Figure S1B) [27,34,35], results in synthetic lethality when combined with the *rad27::LEU2* allele (Figure 1). In previous experiments, we found that *rad27::LEU2* mutant cells display a profusion of DSBs [8]. As both *rad59::LEU2* and *rad59-K166A* substantially reduce association of Rad52 with DSBs [21], we speculate that a critical reduction in the association of Rad52 with the many DSBs in *rad27::LEU2 rad59::LEU2* and *rad27::LEU2 rad59-K166A* double mutants may inhibit their rescue by HR, and results in a lethal level of chromosome loss. The *rad59-F180A* and *rad59-K174A* alleles, which change conserved residues in the same α -helical domain altered by *rad59-K166A*, may have incrementally less severe effects on association of

Table 2 Rates of mutation and unequal sister chromatid recombination in wild-type and mutant haploid strains

Genotype	Mutation rate (10^{-7})	USCR rate (10^{-6})
Wild-type	4.0 (3.8, 7.4) [1]	1.0 (0.8, 1.2) [1]
<i>rad51::LEU2</i>	n.d.	1.4 (1.0, 1.8) [+1.4]
<i>rad59::LEU2</i>	7.5 (6.6, 8.6) [+1.9]	0.82 (0.43, 1.4) [-1.3]
<i>rad59-Y92A</i>	4.4 (3.9, 5.3) [+1.1]	1.3 (1.1, 1.8) [+1.3]
<i>rad59-K174A</i>	3.2 (1.8, 5.5) [-1.3]	1.1 (0.85, 2.1) [+1.1]
<i>rad59-F180A</i>	4.8 (4, 6.9) [+1.2]	0.61 (0.47, 0.95) [-1.6]
<i>rad27::LEU2</i>	200 (90, 590) [+50]	47 (39, 100) [+47]
<i>rad27::LEU2 rad59-Y92A</i>	220 (60, 510) [+55]	39 (25, 99) [+39]
<i>rad27::LEU2 rad59-K174A</i>	130 (110, 190) [+32.5]	38 (33, 53) [+38]
<i>rad27::LEU2 rad59-F180A</i>	190 (110, 500) [+47.5]	60 (49, 120) [+60]

Rates of *CAN1* mutation or USCR were determined from at least 10 independent cultures as described in the Methods. The 95% confidence intervals are in parentheses. Fold decreases (-) and increases (+) from wild-type are in brackets. n.d. - not determined.

Table 3 Rates of loss of heterozygosity in wild-type and mutant diploid strains

Genotype	ILOH rate (10 ⁻⁵)	TLOH rate (10 ⁻⁴)	CL rate (10 ⁻⁵)
Wild-type	2.5 (2.1, 3.1) [1]	0.92 (0.62, 1.2) [1]	3.0 (2.5, 3.9) [1]
<i>rad51::LEU2/rad51::LEU2</i>	1.2 (0.92, 2.5) [-2]	1.3 (0.38, 2) [+1.4]	54 (19, 64) [+18]
<i>rad59::LEU2/rad59::LEU2</i>	1.8 (1.2, 2.9) [-1.4]	1.4 (1.1, 1.9) [+1.5]	6.2 (5.8, 10.2) [+2]
<i>rad59-Y92A/rad59-Y92A</i>	3.2 (2.7, 4.8) [+1.3]	0.95 (0.83, 1.5) [1]	2.5 (2.0, 3.6) [-1.2]
<i>rad59-K174A/rad59-K174A</i>	2.0 (1.3, 3.5) [-1.3]	0.76 (0.40, 1.1) [-1.2]	5.6 (2.9, 8.4) [+1.9]
<i>rad59-F180A/rad59-F180A</i>	3.8 (3.1, 5.1) [+1.5]	0.82 (0.63, 1.7) [-1.1]	3.0 (1.5, 7.9) [1]
<i>rad27::LEU2/rad27::LEU2</i>	28 (25, 64) [+11]	34 (24, 47) [+37]	38 (29, 54) [+13]
<i>rad27::LEU2/rad27::LEU2 rad59-Y92A/rad59-Y92A</i>	28 (13, 56) [+11]	36 (17, 50) [+39]	29 (23, 74) [+9.7]
<i>rad27::LEU2/rad27::LEU2 rad59-K174A/rad59-K174A</i>	26 (22, 55) [+10]	33 (24, 39) [+36]	32 (18, 48) [+11]
<i>rad27::LEU2/rad27::LEU2 rad59-F180A/rad59-F180A</i>	52 (29, 76) [+21]	35 (22, 57) [+38]	57 (18,124) [+19]

Rates of interstitial LOH (ILOH), terminal LOH (TLOH), and chromosome loss (CL) from a minimum of 12 independent cultures were determined as described in the Methods. The 95% confidence intervals are in parentheses. Fold decreases (-) and increases (+) from wild-type are in brackets.

Rad52 with DSBs. This may result in their serially reduced inhibition of repair of replication-induced DSBs by HR (Figure 3C; Additional file 1: Table S2) and commensurate effects on growth (Table 1; Additional file 1: Table S2) when combined with *rad27*. An accumulation of *rad27::LEU2 rad59-F180A* double mutant cells in the G2 phase of the cell cycle, as compared to *rad27::LEU2* single mutant or *rad27::LEU2 rad59-K174A* double mutant cells is consistent with more deficient repair of replication-induced DSBs by HR (Figure 3). This further supports the notion that *RAD59* promotes the survival of *rad27::LEU2* mutant cells by facilitating the rescue of replication lesions by HR. Recently, *RAD59* has been shown to be required for the viability of DNA ligase I-deficient mutants, verifying the requirement for this factor in accommodating to incomplete DNA replication [51].

In striking contrast to the other *rad59* alleles, *rad59-Y92A* stimulated HR (Figure 3B; Figure 4B). This hyper-recombinogenic effect was distinct from that caused by *rad27* as it was not accompanied by significant effects on doubling time (Table 1), cell cycle profile (Figure 2), mutation (Table 2), unequal sister chromatid exchange, or LOH (Table 3), suggesting that *rad59-Y92A* does not cause an accumulation of replication lesions. The observation that the stimulatory effect of *rad59-Y92A* was completely suppressed by a null allele of *RAD51*, and was mutually epistatic with a null allele of *SRS2* (Figure 3D), suggests that *rad59-Y92A* may increase HR by increasing the stability of Rad51-DNA filaments, perhaps by changing its interaction with Rad51 (24). An increase in DSBs combined with an increase in the stability of Rad51 filaments at the DSBs may underlay the synergistically increased rates of HR observed in *rad27 rad59-Y92A* double mutants (Figures 3C and 4B). However, since Rad59 also interacts with RPA [52] and RSC [53], the increase in HR observed in *rad59-Y92A* mutant cells may also involve changes in additional processes.

While our results support a prominent role for *RAD59*-dependent HR in the repair of replication lesions in *rad27::LEU2* mutants, HR mechanisms that do not depend on *RAD59* were also strongly stimulated in *rad27::LEU2* mutants. In particular, inter-chromosomal HR leading to interstitial and terminal LOH events was strongly stimulated by *rad27*, but was unaffected by the *rad59* alleles (Table 3). Inter-chromosomal HR leading to LOH is thought to occur by break-induced replication (BIR) [54]. BIR has been proposed to utilize a single-ended DSB on one homolog to generate a replication fork-like intermediate with the unbroken homolog that may potentially proceed until reaching the end of the donor chromosome (Additional file 1: Figure S4A) [22]. In contrast, *RAD59*-dependent heteroallelic recombination is thought to utilize a double-ended DSB where both ends are rescued, either through concerted interactions with the unbroken homolog, or through the first end interacting with the homolog followed by the second end annealing with the first after gaining sequences copied from the unbroken homolog (Additional; file 1: Figure S4B). The stimulation of both mechanisms of HR between homologs suggests that loss of *RAD27* leads to the accumulation of both single- and double-ended DSBs. DSBs may arise when the failure to remove flaps on the 5' ends of Okazaki fragments leads to accumulation of nicks on newly replicated lagging strands (Figure 5). Persistence of these nicks into the subsequent cell cycle will leave discontinuities on the template for leading strand synthesis that will stall replication and form single-ended DSBs. If a second replication fork from an adjacent replicon collides with the first stalled fork, a double-ended DSB can arise. A genome-wide increase in replication-induced DSB formation, like that induced by many chemotherapeutic agents, would therefore require a robust response by the HR apparatus to prevent chromosome loss, potentially explaining the critical role of HR in determining sensitivity to these drugs in humans [55,56].




Figure 5 Models for initiation of RAD51- and RAD59-dependent and -independent HR by defective lagging strand synthesis.

1.) Accumulation of daughter strand nicks in the absence of Rad27 nuclease causes replication fork stalling during the next S phase when the lagging strand becomes the template for leading strand synthesis and the replication fork encounters the discontinuity. 2.) The stalled fork is converted into an intact chromatid and a single-ended DSB. The single-ended DSB becomes a substrate for RAD51- and RAD59-independent HR mechanisms, such as interstitial and terminal LOH (Additional file 1: Figure S3). 3.) The replication fork from an upstream replicon converges with the previously stalled fork. 4.) Converged forks are converted into an intact chromatid and a double-ended DSB. The double-ended DSB becomes a substrate for RAD51- and RAD59-dependent HR mechanisms, such as ectopic gene conversion and heteroallelic recombination (Figures 3A and 4A).

Conclusions

RAD59 encodes one of several homologous recombination (HR) factors required for viability of budding yeast cells lacking the DNA replication factor, Rad27. This demonstrates that the HR apparatus is required to prevent the lethal effects of dysfunctional replication, but no link between a specific HR mechanism and survival has been previously established. In this analysis, we show that two *rad59* alleles that diminish association of Rad52 with double-strand breaks are synthetically lethal with *rad27*, while two others coordinately reduce RAD51-dependent HR and growth, thus linking RAD51-dependent repair with survival. Another allele stimulates HR by stabilizing Rad51-DNA filaments. Therefore, Rad59 influences the repair of replication lesions by HR through its interactions with multiple HR factors. We speculate that the massive increase in replication failure genome-wide that results from loss of Rad27 may be similar to that caused by chemotherapeutic agents in human cells, potentially explaining why the HR apparatus is critical in determining sensitivity to these drugs.

Methods

Strains

All strains used in this study were isogenic and are listed in Additional file 1: Table S1. Standard techniques for yeast strain construction and growth were used [57]. Construction of the *rad27::LEU2*, *rad51::LEU2*, *rad59::*

LEU2, *rad59-Y92A*, *rad59-K166A*, *rad59-K174A*, *rad59-F180A* and *srs2::TRP1* alleles have been described previously [27,58-60]. The *rad27::LEU2* allele can be followed in crosses by PCR, using the forward primer 5'-GCG TTG ACA GCA TAC ATT-3', and reverse primer 5'-CGT ACA AAC CAA ATG CGG-3'. The *rad59::LEU2* allele is followed by PCR using the forward primer 5'-GCC ACA GTT TGG CAA GGG-3', and the reverse primer 5'-GGG TTT GTT GCC ATC TGC G-3'. The *rad59* missense alleles were followed in crosses by allele-specific PCR [27]. Unique forward primers were used to detect *rad59-Y92A* (5'-GCT AAT GAA ACA TTC GGG GC-3'), *rad59-K166A* (5'-AAT GTT ATA ACA GGT CGA AAG C-3'), *rad59-K174A* (5'-AAG GGT TAC GTA GAG GAG AAG-3'), and *rad59-F180A* (5'-AAG AAG GCG TTA TTG AGC GC-3'). All allele-specific PCRs use the same reverse primer (5'-TAT ATA AGT ACG TGA GAT CTA TTT G-3'). Presence of the *rad59-K174A* allele is scored by digesting the PCR product with *MseI* restriction endonuclease. DNA was purified for PCR analysis using a standard method [61].

Synthetic lethality

Diploid yeast strains heterozygous for each of the *rad59* alleles (*rad59/RAD59*) and the *rad27::LEU2* allele (*rad27::LEU2/RAD27*) were sporulated and dissected. After 72 h, five representative tetrads from each diploid were selected. The presence of *rad27* and *rad59* mutant alleles in each of

the colonies that arose from the spores was scored using PCR as described above.

Doubling time

At least 10, five-milliliter YPD (1% yeast extract, 2% peptone, 2% dextrose) cultures were inoculated with colonies arising from the spores of freshly dissected tetrads and grown overnight at 30°. These were sub-cultured into Klett tubes containing five milliliters of YPD medium that were incubated at 30° while shaking. Cell density was measured by monitoring culture turbidity with a Klett-Summerson colorimeter each hour over a 10 h period. Doubling times were calculated using a standard algorithm [62]. The 95% confidence intervals and Mann–Whitney values were determined using the Prism statistics package (GraphPad, La Jolla, CA).

Flow cytometry

At least five, five-milliliter YPD cultures were inoculated with colonies arising from freshly dissected tetrads and grown overnight at 30°. Overnight cultures were sub-cultured into five milliliters of YPD medium and grown to mid-log phase at 30° defined by growth curve using a Klett-Summerson colorimeter. Cells were processed for flow cytometry using the following adaptation of a published method [63]. The cell density was determined by hemacytometer count and aliquots containing 10^7 cells were pelleted, resuspended in 70% ice-cold ethanol, and fixed while rotating at 4° overnight. Fixed cells were pelleted, resuspended in 1 ml of citrate buffer (50 mM Na citrate, pH 7.2), and sonicated (Misonix 3000, Farmingdale, NY). Sonicated cells were pelleted, resuspended in citrate buffer and treated with 25 μ l of 10 mg/ml RNase A, at 50° for one h, followed by treatment with 50 μ l of 20 mg/ml Proteinase K and incubation at 50° for one h. Cells were pelleted and resuspended in 1 ml of citrate buffer, and either rotated overnight at 4°, or stained immediately by adding 16 μ l of 1 mg/ml propidium iodide and rotating for 45 min at room temperature in the dark before processing by flow cytometry (Beckman Coulter CyAn ADP 9color, Miami FL). Fractions of cells in the G1, S and G2/M phases of the cell cycle were determined using FlowJo v.7.6.5 image processing software (Tree Star, Ashland, OR). The ratio of cells in G1 vs. S + G2/M were calculated for each trial and the median value for each strain used for comparing cell cycle distributions in different strains. The Mann–Whitney test was used to assess the statistical significance of differences between strains.

Spontaneous ectopic gene conversion

Spontaneous ectopic gene conversion in haploid strains was assayed as described previously [64], but using substrates described in a separate analysis [41]. All strains

contained the *sam1- Δ Bgl II-HOcs* allele at the *SAM1* locus on chromosome XII, the *sam1- Δ Sal I* allele adjacent to the *HIS3* locus on chromosome XV, and a *HIS3* gene replacing the *SAM2* coding sequence at the *SAM2* locus (*sam2::HIS3*) on chromosome IV. The *sam1- Δ Bgl II-HOcs* allele has a 117 bp fragment of the *MAT* locus disrupting the *Bgl II* site in the *SAM1* coding sequence, while the *sam1- Δ Sal I* allele has a 4 bp insertion at the *Sal I* site [41]. The *sam1- Δ Sal I* allele lacks a promoter, preventing conversion events at this locus from generating AdoMet⁺ recombinants. The *sam1- Δ Bgl II-HOcs* and *sam1- Δ Sal I* alleles are also in opposite orientations relative to their centromeres, preventing the isolation of single crossover recombinants.

At least ten freshly dissected haploid segregants of each strain were inoculated into five-milliliter YPD cultures supplemented with 100 μ g/ml of *S*-adenosylmethionine (AdoMet) and grown to saturation at 30°. Appropriate dilutions of each culture were plated onto YPD + AdoMet plates to determine the number of viable cells, and onto YPD plates lacking AdoMet to determine the number of AdoMet prototrophic recombinants.

All rates were determined by the method of the median [65]. Rates and 95% confidence intervals were calculated as described previously [66].

Spontaneous hetero-allelic recombination

Rates of spontaneous hetero-allelic recombination were determined as for ectopic gene conversion except that different substrates were used in diploid cells. All strains contained the *sam2- Δ EcoR V-HOcs* allele at the *SAM2* locus on one copy of chromosome IV, the *sam2- Δ Sal I* allele on the other, and a *LEU2* marker replacing the *SAM1* coding sequence at the *SAM1* locus on both copies of chromosome XII. The *sam2- Δ EcoR V-HOcs* allele has a 117 bp fragment of the *MAT* locus disrupting the *EcoR V* site, while the *sam2- Δ Sal I* allele has a 4 bp insertion disrupting the *Sal I* site [41].

Mutation rate

Rates of mutation at the *CAN1* locus were examined using a previously published assay [8,10,18]. At least ten freshly dissected segregants were used to inoculate one-milliliter YPD cultures that were grown to saturation at 30°. Appropriate dilutions were plated onto YPD to determine viability and synthetic medium lacking arginine but containing 60 μ g/ml of canavanine to select for mutants.

Unequal sister chromatid recombination (USCR)

Rates of USCR were determined using a previously published assay [8,10,67]. At least ten freshly dissected segregants containing the USCE construct at the *TRP1* locus on chromosome IV and the *his3 Δ 200* allele at the

HIS3 locus on chromosome XV, were struck out to single colonies on YPD. After three days of growth at 30°, single colonies were used to inoculate one-milliliter YPD cultures, and grown to saturation at 30°. Appropriate dilutions were plated onto YPD to assess viability and onto medium lacking histidine to determine the number of histidine prototrophic recombinants.

Loss of heterozygosity (LOH)

Rates of spontaneous LOH by three different mechanisms were assessed using a previously published assay [8]. Freshly dissected haploid segregants containing either the *hxt13::URA3*, *CAN1*, and *HOM3* alleles, or the *HXT13*, *can1-100*, and *hom3-10* alleles on chromosome V were crossed and the resulting diploids struck out to single colonies on YPD. At least 12 independent colonies were inoculated into one-milliliter YPD liquid cultures and grown to saturation at 30°. Appropriate dilutions were plated onto YPD for viability and synthetic medium lacking arginine, but containing 60 µg/ml of canavanine to select for clones resistant to canavanine. After three days of growth at 30° canavanine-resistant (Can^{R}) colonies were replica plated onto synthetic medium lacking either uracil or threonine to assay for the presence of the *hxt13::URA3* (Ura^+) and *HOM3* (Thr^+) alleles, respectively. Rates of interstitial LOH, terminal LOH, and CL were determined from the number of $\text{Ura}^+\text{Can}^{\text{R}}\text{Thr}^+$, $\text{Ura}^-\text{Can}^{\text{R}}\text{Thr}^+$, and $\text{Ura}^-\text{Can}^{\text{R}}\text{Thr}^-$ recombinant colonies, respectively.

Modeling the Rad59 protein

The crystal structure of the N-terminus of human Rad52 [34] was obtained from the RCSB Protein Data Bank (<http://www.rcsb.org/pdb/>). This structure was imaged using the molecular modeling program, SYBYL, and the amino acids corresponding to those mutated in the *rad59* missense alleles were identified, and highlighted.

Availability of supporting data

The data sets supporting the results of this article are included within the article and in Additional file 1.

Additional file

Additional file 1: Table S1. *Saccharomyces cerevisiae* strains used in this study. **Table S2.** Summary of quantitative data. **Figure S1. A.** Multiple amino acid sequence alignment of ScRad59 with ScRad52 and HsRad52. **B.** Molecular modeling of the proteins encoded by the *rad59* missense alleles demonstrates that Rad59-Y92A is in a different structural motif. **Figure S2.** The unequal sister chromatid recombination (USCR) assay for measuring spontaneous homologous recombination between sister chromatids in haploid yeast. **Figure S3.** The loss of heterozygosity assay for measuring spontaneous Rad51-independent homologous recombination. **Figure S4.** LOH is the recombination product of a single-ended DSB, whereas HAR results from repair of a double-ended DSB. **A)** LOH results from the repair of a single-ended DSB by HR. **B)** HAR results from the repair of a double-ended DSB by HR.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

LL carried out the synthetic lethality experiments, LOH genetic studies, flow cytometric analysis, sequence alignment, designed the figures and tables, and drafted the manuscript. GM performed the growth, mutation and USCR rate studies. SO assisted with the synthetic lethality and LOH experiments. BF contributed to the LOH experiments. AB conceived of the study, designed and carried out the ectopic gene conversion and hetero-allelic recombination analyses, and helped draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank M. Boldin, M. Kalkum, R.-J. Lin, T. O'Connor, and J. Stark for stimulating discussions, and N. Pannunzio for comments on the manuscript. We would like to acknowledge the City of Hope Biostatistics and Bioinformatics, and Flow Cytometry Core Facilities for their assistance. This work was supported by a Morgan and Helen Chu graduate student fellowship to L.C.L., a summer undergraduate research fellowship from the Howard Hughes Medical Institute to S.N.O., a summer student fellowship from the Eugene and Ruth Roberts Summer Academy to B.X.H.F., and funds from the Beckman Research Institute of the City of Hope.

Author details

¹Department of Molecular and Cellular Biology, Beckman Research Institute of the City of Hope, 91010 Duarte, CA, USA. ²Irell & Manella Graduate School of Biological Sciences, Beckman Research Institute of the City of Hope, 91010 Duarte, CA, USA. ³Occidental College, 90041 Los Angeles, CA, USA. ⁴Department of Genetics, Stanford University School of Medicine, 94305 Stanford, CA, USA.

Received: 22 August 2013 Accepted: 4 October 2013

Published: 14 October 2013

References

- Gordenin DA, Malkova AL, Peterzen A, Kulikov VN, Pavlov YI, Perkins E, Resnick MA: Transposon Tn5 excision in yeast: Influence of DNA polymerases α , δ , and ϵ and repair genes. *Proc Natl Acad Sci U S A* 1992, **89**:3785–3789.
- Vallen EA, Cross FR: Mutations in *RAD27* define a potential link between G1 cyclins and DNA replication. *Mol Cell Biol* 1995, **15**(8):4291–4302.
- Ruskin B, Fink G: Mutations in *POL1* increase the mitotic instability of tandem inverted repeats in *Saccharomyces cerevisiae*. *Genetics* 1993, **133**:43–56.
- Tishkoff DX, Boerger AL, Bertrand P, Filosi N, Gaida GM, Kane MF, Kolodner RD: Identification and characterization of *Saccharomyces cerevisiae* *EXO1*, a gene encoding an exonuclease that interacts with *MSH2*. *Proc Natl Acad Sci U S A* 1997, **94**:7487–7492.
- Zou H, Rothstein R: Holliday junctions accumulate in replication mutants via a RecA homolog-independent mechanism. *Cell* 1997, **90**:87–96.
- Chen C, Kolodner R: Gross chromosomal rearrangements in *Saccharomyces cerevisiae* replication and recombination defective mutants. *Nat Genet* 1999, **23**:81–85.
- Pavlov YI, Shcherbakova PV, Kunkel TA: *In vivo* consequences of putative active site mutations in yeast DNA Polymerases α , ϵ , δ , and ζ . *Genetics* 2001, **159**:47–64.
- Navarro MS, Bi L, Bailis AM: A mutant allele of the transcription factor IIIH helicase gene, *RAD3*, promotes loss of heterozygosity in response to a DNA replication defect in *Saccharomyces cerevisiae*. *Genetics* 2007, **176**:1391–1402.
- Venkatesan RN, Treuting PM, Fuller ED, Goldsby RE, Norwood TH, Gooley TA, Ladiges WC, Preston BD, Loeb LA: Mutation at the polymerase active site of mouse DNA polymerase δ increases genomic instability and accelerates tumorigenesis. *Mol Cell Biol* 2007, **27**(21):7669–7682.
- Mito E, Mokhnatkin JV, Steele MC, Buettner VL, Sommer SS, Manthey GM, Bailis AM: Mutagenic and recombinogenic responses to defective DNA polymerase δ are facilitated by the Rev1 protein in *pol3-t* mutants of *Saccharomyces cerevisiae*. *Genetics* 2008, **179**:1795–1806.

11. Galli A, Cervelli T, Schiestl RH: Characterization of the hyperrecombination phenotype of the *pol3-t* mutation of *Saccharomyces cerevisiae*. *Genetics* 2003, **164**:65–79.
12. Harrington JJ, Lieber MR: The characterization of a mammalian DNA structure-specific endonuclease. *EMBO J* 1994, **13**(5):1235–1246.
13. Liu Y, Kao HI, Bambara RA: Flap endonuclease 1: A central component of DNA metabolism. *Annu Rev Biochem* 2004, **73**:589–615.
14. Wu X, Wang Z: Relationships between yeast Rad27 and Apn1 in response to apurinic/aprimidinic (AP) sites in DNA. *Nucleic Acids Res* 1999, **27**(4):956–962.
15. Tseng HM, Tomkinson AE: Processing and joining of DNA ends coordinated by interactions among Dnl4/Lif1, Pol4, and FEN-1. *J Biol Chem* 2004, **279**(46):47580–47588.
16. Parenteau J, Wellinger RJ: Accumulation of single-stranded DNA and destabilization of telomeric repeats in yeast mutant strains carrying a deletion of *RAD27*. *Mol Cell Biol* 1999, **19**(6):4143–4152.
17. Nowosielska A: Bacterial DNA repair genes and their eukaryotic homologues: 5. The role of recombination in DNA repair and genome stability. *Acta Biochimica Polonica* 2007, **54**(3):483–494.
18. Tishkoff DX, Filosi N, Gaida GM, Kolodner RD: A novel mutation avoidance mechanism dependent on *S. cerevisiae* *RAD27* is distinct from DNA mismatch repair. *Cell* 1997, **88**:253–263.
19. Symington LS: Homologous recombination is required for the viability of *rad27* mutants. *Nucleic Acids Res* 1998, **26**(24):5589–5595.
20. Debrauwere H, Loeillet S, Lin W, Lopes J, Nicolas A: Links between replication and recombination in *Saccharomyces cerevisiae*: a hypersensitive requirement for homologous recombination in the absence of Rad27 activity. *Proc Natl Acad Sci U S A* 2001, **98**(15):8263–8269.
21. Pannunzio NR, Manthey GM, Liddell LC, Fu BX, Roberts CM, Bailis AM: Rad59 regulates association of Rad52 with DNA double-strand breaks. *Microbiology Open* 2012, **1**(3):285–297.
22. Paques F, Haber JE: Multiple pathways of recombination induced by double-strand breaks in *Saccharomyces cerevisiae*. *Micro Mol Biol Rev* 1999, **63**(2):349–404.
23. Krogh BO, Symington LS: Recombination proteins in yeast. *Annu Rev Genet* 2004, **38**:233–271.
24. Wu Y, Kantake N, Sugiyama T, Kowalczykowski SC: Rad51 protein controls Rad52-mediated DNA annealing. *J Biol Chem* 2008, **283**(21):14883–14892.
25. Davis AP, Symington LS: The yeast recombinational repair protein Rad59 interacts with Rad52 and stimulates single-strand annealing. *Genetics* 2001, **159**:515–525.
26. Pannunzio NR, Manthey GM, Bailis AM: *RAD59* is required for efficient repair of simultaneous double-strand breaks resulting in translocations in *Saccharomyces cerevisiae*. *DNA Repair (Amst)* 2008, **7**(5):788–800.
27. Pannunzio NR, Manthey GM, Bailis AM: Rad59 and Rad1 cooperate in translocation formation by single-strand annealing in *Saccharomyces cerevisiae*. *Curr Genet* 2010, **56**(1):87–100.
28. Sugawara N, Ira G, Haber JE: DNA length dependence of the single-strand annealing pathway and the role of *Saccharomyces cerevisiae* *RAD59* in double-strand break repair. *Mol Cell Biol* 2000, **20**(14):5300–5309.
29. Bai Y, Symington LS: A Rad52 homolog is required for *RAD51*-independent mitotic recombination in *Saccharomyces cerevisiae*. *Genes Dev* 1996, **10**(16):2025–2037.
30. Cortes-Ledesma F, Tous C, Aguilera A: Different genetic requirements for repair of replication-born double-strand breaks by sister-chromatid recombination and break-induced replication. *Nucleic Acids Res* 2007, **35**(19):6560–6570.
31. Mott C, Symington LS: *RAD51*-independent inverted-repeat recombination by a strand-annealing mechanism. *DNA Repair (Amst)* 2011, **10**(4):408–415.
32. Cortes-Ledesma F, Malagon F, Aguilera A: A novel yeast mutation, *rad52-L89F*, causes a specific defect in Rad51-independent recombination that correlates with a reduced ability of *Rad52-L89F* to interact with Rad59. *Genetics* 2004, **168**:553–557.
33. Feng Q, Doring L, de Mayolo AA, Lettier G, Lisby M, Erdeniz N, Mortensen UH, Rothstein R: Rad52 and Rad59 exhibit both overlapping and distinct functions. *DNA Repair (Amst)* 2007, **6**(1):27–37.
34. Kagawa W, Kurumizaka H, Ishitani R, Fukai S, Nureki O, Shibata T, Yokoyama S: Crystal structure of the homologous-pairing domain from the human Rad52 recombinase in the undecameric form. *Mol Cell* 2002, **10**:359–371.
35. Lloyd JA, McGrew DA, Knight KL: Identification of residues important for DNA binding in the full-length human Rad52 protein. *J Mol Biol* 2005, **345**(2):239–249.
36. Veaute X, Jeusset J, Soustelle C, Kowalczykowski SC, Fabre F: The Srs2 helicase prevents recombination by disrupting Rad51 nucleoprotein filaments. *Nature* 2003, **423**:309–312.
37. Antony E, Tomko EJ, Xiao Q, Krejci L, Lohman TM, Ellenberger T: Srs2 disassembles Rad51 filaments by a protein-protein interaction triggering ATP turnover and dissociation of Rad51 from DNA. *Mol Cell* 2009, **35**(1):105–115.
38. Sung P: Catalysis of ATP-dependent homologous DNA pairing and strand exchange by yeast RAD51 protein. *Science* 1994, **265**(5176):1241–1243.
39. Bai Y, Davis AP, Symington LS: A novel allele of *RAD52* that causes severe DNA repair and recombination deficiencies only in the absence of *RAD51* or *RAD59*. *Genetics* 1999, **153**:1117–1130.
40. Jablonovich Z, Liefshitz B, Steinlauf R, Kupiec M: Characterization of the role played by the *RAD59* gene of *Saccharomyces cerevisiae* in ectopic recombination. *Curr Genet* 1999, **36**:13–20.
41. Bailis AM, Maines S, Negritto MT: The essential helicase gene *RAD3* suppresses short-sequence recombination in *Saccharomyces cerevisiae*. *Mol Cell Biol* 1995, **15**(5):3998–4008.
42. Liefshitz B, Parket A, Maya R, Kupiec M: The role of DNA repair genes in recombination between repeated sequences in yeast. *Genetics* 1995, **140**:1199–1211.
43. Rong L, Klein HL: Purification and characterization of the SRS2 DNA helicase of the yeast *Saccharomyces cerevisiae*. *J Biol Chem* 1993, **268**(2):1252–1259.
44. Rong L, Palladino F, Aguilera A, Klein HL: The hyper-gene conversion *hpr5-1* mutation of *Saccharomyces cerevisiae* is an allele of the *SRS2/RADH* gene. *Genetics* 1991, **127**:75–85.
45. Palladino F, Klein HL: Analysis of mitotic and meiotic defects in *Saccharomyces cerevisiae* *SRS2* DNA helicase mutants. *Genetics* 1992, **132**(1):23–37.
46. Morrison DP, Hastings PJ: Characterization of the mutator mutation *mut5-1*. *Mol Gen Genet* 1979, **175**(1):57–65.
47. Lopes J, Ribeyre C, Nicolas A: Complex minisatellite rearrangements generated in the total or partial absence of Rad27/hFEN1 activity occur in a single generation and are Rad51 and Rad52 dependent. *Mol Cell Biol* 2006, **26**(17):6675–6689.
48. Freudenreich CH, Kantrow SM, Zakian VA: Expansion and length-dependent fragility of CTG repeats in yeast. *Science* 1998, **279**(853):853–856.
49. Johnson RE, Kovvali GK, Prakash L, Prakash S: Role of yeast Rth1 nuclease and its homologs in mutation avoidance, DNA repair, and DNA replication. *Curr Genet* 1998, **34**:21–29.
50. Fasullo MT, Davis RW: Direction of chromosome rearrangements in *Saccharomyces cerevisiae* by use of *his3* recombinational substrates. *Mol Cell Biol* 1988, **8**(10):4370–4380.
51. Nguyen HD, Becker J, Thu YM, Costanzo M, Koch EN, Smith S, Myers CL, Boone C, Bielsky AK: Unligated Okazaki fragments induce PCNA ubiquitination and a requirement for Rad59-dependent replication fork progression. *PLoS One* 2013, **8**(6):e6379.
52. Davis AP, Symington LS: The Rad52-Rad59 complex interacts with Rad51 and replication protein A. *DNA Repair (Amst)* 2003, **2**:1127–1134.
53. Oum J-H, Seong C, Kwon Y, Ji J-H, Sid A, Ramakrishnan S, Ira G, Malkova A, Sung P, Lee SE, Shim EY: RSC facilitates Rad59-dependent homologous recombination between sister chromatids by promoting cohesin loading at DNA double-strand breaks. *Mol Cell Biol* 2011, **31**(19):3924–3937.
54. Pohl TJ, Nickoloff JA: Rad51-independent interchromosomal double-strand break repair by gene conversion requires Rad52 but not Rad55, Rad57, or Dmc1. *Mol Cell Biol* 2008, **28**(3):897–906.
55. Nikolova T, Ensminger M, Lobrich M, Kaina B: Homologous recombination protects mammalian cells from replication-associated DNA double-strand breaks arising in response to methyl methanesulfonate. *DNA Repair (Amst)* 2010, **9**(10):1050–1063.
56. Nikolova T, Hennekes F, Bhatti A, Kaina B: Chloroethylnitrosourea-induced cell death and genotoxicity: cell cycle dependence and the role of DNA double-strand breaks. *HR and NHEJ. Cell Cycle* 2012, **11**(14):2606–2619.
57. Sherman F, Fink F, Hicks J: *Methods in Yeast Genetics*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 1986.
58. Schild D, Konforti B, Perez C, Gish W, Mortimer RK: Isolation and characterization of yeast DNA repair genes. I. Cloning of the *RAD52* gene. *Curr Genet* 1983, **7**:85–92.
59. Schild D, Calderon IL, Contopoulos R, Mortimer RK: *Cloning of yeast recombination repair genes and evidence that several are nonessential genes*. New York: Alan R. Liss; 1983.

60. Frank G, Qiu J, Somsouk M, Weng Y, Somsouk L, Nolan JP, Shen B: **Partial functional deficiency of E160D flap endonuclease-1 mutant *in vitro* and *in vivo* is due to defective cleavage of DNA substrates.** *J Biol Chem* 1998, **273**(49):33064–33072.
61. Hoffman CS, Winston F: **A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*.** *Gene* 1987, **57**(2–3):267–272.
62. Singleton P: *Bacteria in Biology, Biotechnology, and Medicine*. New York: John Wiley & Sons; 1995.
63. Nash N, Tokiwa G, Anand S, Erickson K, Fitcher AB: **The *WHI1+* gene of *Saccharomyces cerevisiae* tethers cell division to cell size and is a cyclin homolog.** *EMBO J* 1988, **7**(13):4335–4346.
64. Bailis AM, Rothstein R: **A defect in mismatch repair in *Saccharomyces cerevisiae* stimulates ectopic recombination between homeologous genes by an excision repair dependent process.** *Genetics* 1990, **126**:535–547.
65. Lea DE, Coulson CA: **The distribution of the numbers of mutants in bacterial populations.** *J Genet* 1949, **49**:264–285.
66. Spell RM, Jinks-Robertson S: **Determination of mitotic recombination rates by fluctuation analysis in *Saccharomyces cerevisiae*.** *Methods Mol Biol* 2004, **262**:3–12.
67. Fasullo MT, Davis RW: **Recombinational substrates designed to study recombination between unique and repetitive sequence *in vivo*.** *Proc Natl Acad Sci U S A* 1987, **84**:6215–6219.

doi:10.1186/1471-2180-13-229

Cite this article as: Liddell *et al.*: Alleles of the homologous recombination gene, *RAD59*, identify multiple responses to disrupted DNA replication in *Saccharomyces cerevisiae*. *BMC Microbiology* 2013 **13**:229.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

