Research article

Open Access

Hfq regulates the expression of the thermostable direct hemolysin gene in *Vibrio parahaemolyticus*

Masayuki Nakano, Akira Takahashi*, Zehong Su, Nagakatsu Harada, Kazuaki Mawatari and Yutaka Nakaya

Address: Department of Nutrition and Metabolism, Institute of Health Biosciences, The University of Tokushima Graduate School, Tokushima, Japan

Email: Masayuki Nakano - mnakano@nutr.med.tokushima-u.ac.jp; Akira Takahashi* - akiratak@nutr.med.tokushima-u.ac.jp; Zehong Su - zehongsu@hotmail.com; Nagakatsu Harada - harada@nutr.med.tokushima-u.ac.jp; Kazuaki Mawatari - mawatari@nutr.med.tokushima-u.ac.jp; Yutaka Nakaya - yutaka-nakaya@nutr.med.tokushima-u.ac.jp

* Corresponding author

Published: 21 September 2008

Received: 4 February 2008

BMC Microbiology 2008, 8:155 doi:10.1186/1471-2180-8-155

Accepted: 21 September 2008

This article is available from: <http://www.biomedcentral.com/1471-2180/8/155>

© 2008 Nakano et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The *hfq* gene is conserved in a wide variety of bacteria and Hfq is involved in many cellular functions such as stress responses and the regulation of gene expression. It has also been reported that Hfq is involved in bacterial pathogenicity. However, it is not clear whether Hfq regulates virulence in *Vibrio parahaemolyticus*. To evaluate this, we investigated the effect of Hfq on the expression of virulence-associated genes including thermostable direct hemolysin (TDH), which is considered to be an important virulence factor in *V. parahaemolyticus*, using an *hfq* deletion mutant.

Results: The production of TDH in the *hfq* deletion mutant was much higher than in the parental strain. Quantification of *tdh* promoter activity and mRNA demonstrated that transcription of the *tdh* gene was up-regulated in the mutant strain. The *hfq*-complemented strain had a normal (parental) amount of *tdh* expression. The transcriptional activity of *tdhA* was particularly increased in the mutant strain. These results indicate that Hfq is closely associated with the expression level of the *tdh* gene. Interestingly, other genes involved in the pathogenicity of *V. parahaemolyticus*, such as VPI680, *vopC*, and *vopT*, were also up-regulated in the mutant strain.

Conclusion: Hfq regulates the expression of virulence-associated factors such as TDH and may be involved in the pathogenicity of *V. parahaemolyticus*.

Background

Vibrio parahaemolyticus is a major causative agent of gastroenteritis, and is often associated with the consumption of raw or undercooked shellfish. In addition to gastroenteritis, this organism causes clinical manifestations such as wound infections and septicemia [1-3]. Most clinical isolates of *V. parahaemolyticus* exhibit β -type hemolysis on a special blood agar medium, whereas environmental iso-

lates do not. This hemolysis is called the Kanagawa phenomenon (KP) and is considered to be a reliable marker for detection and identification of pathogenic strains [4]. It has been reported that thermostable direct hemolysin (TDH) specifically contributes to the KP [2]. Previous studies have shown that TDH shows several biological activities such as hemolysis, cell cytotoxicity and enterotoxicity [5-7]. Furthermore, molecular epidemiological

studies have indicated that the *tdh* gene is strongly associated with clinical strains [8]. Therefore, TDH is thought to be an important virulence factor in *V. parahaemolyticus*. Although the genetic structures of the *tdh* gene and its neighboring regions are known, there is little information as to how *tdh* gene expression is controlled by other factors in this organism.

The *hfq* gene is conserved in a wide range of bacteria, but Hfq varies in the number of amino acids between different bacteria [9]. Hfq is a global regulator that modulates the stability and translation of mRNAs. It is thought that Hfq acts as an RNA chaperone by mediating interactions between many small regulatory RNAs and their mRNA targets [9]. It has been shown that Hfq regulates the expression of many genes, including the genes encoding for stationary-phase sigma factor (*rpoS*) and outer membrane protein A (*ompA*) [10-12]. It has also been shown that Hfq contributes to pathogenicity in bacteria such as *Brucella abortus*, *Listeria monocytogenes*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*, *V. cholerae*, and *Yersinia enterocolitica* [13-18]. Although Hfq is associated with bacterial virulence, it is not clear whether Hfq is associated with pathogenicity in *V. parahaemolyticus*.

In this study, we have attempted to elucidate the correlation between Hfq and *V. parahaemolyticus* pathogenicity, particularly with regard to the important virulence factor TDH, using an *hfq* gene-deletion mutant.

Results

Effect of Hfq on growth of *V. parahaemolyticus*

The complete genome of *V. parahaemolyticus* strain RIMD2210633 has been sequenced, and the *hfq* gene is located on chromosome 1 (GenBank: [BA000031](#)) [19]. The gene structures neighboring *hfq* in *V. parahaemolyticus* are identical to those of *V. cholerae* and *Escherichia coli* (Fig. 1A) [14], and the Hfq amino acid sequence of *V. parahaemolyticus* is highly conserved with that of *V. cholerae* (Fig. 1B).

Previous studies showed that defects in the *hfq* gene decreased the growth rates of *E. coli* and *P. aeruginosa*, but not of *S. typhimurium* and *V. cholerae* [11,14,15,20]. In LB medium supplemented with 3% NaCl, Δhfq , a derivative of the *V. parahaemolyticus* strain RIMD2210633 with an in-frame deletion of 85 of the 87 amino acid residues of Hfq, showed a reduced growth rate as compared with that of the parental strain (Fig. 2). This indicates that Hfq has an effect on the growth of *V. parahaemolyticus*.

Hemolytic activity of secreted protein from Δhfq

Hfq regulates the bacterial pathogenicity that is mediated by alterations in virulence gene expression [13-18]. Therefore, we investigated the hemolytic activity of the Δhfq

strain by determining the level of TDH in culture supernatants, because TDH is thought to be important for *V. parahaemolyticus* pathogenicity. In the following experiments, we evaluated all of the strains under the same growth phase conditions. The relative hemolytic activity in the culture supernatant of the Δhfq strain was significantly higher than that of the parental strain. We introduced pHfq, which harbors the *hfq* gene, into Δhfq and showed that the relative hemolytic activity in the culture supernatant of pHfq/ Δhfq was lower than that of Δhfq . The hemolytic activity in the culture supernatant of pSN1/ Δhfq , in which an empty vector was introduced into Δhfq , was almost the same as that of Δhfq (Fig. 3). These results suggest that Hfq is associated with the production of TDH in *V. parahaemolyticus*.

Hfq is involved in the expression of TDH

We next investigated the level of TDH by Western blotting, and found that the expression of TDH was increased at mid-log growth phase ($OD_{600} = 0.6$) as compared with the parental strain (Fig. 4A). The amount of TDH in the Δhfq strain at stationary growth phase was also higher than that of the parent strain (Fig. 4A). The elevation in TDH production was fully reversed by trans-complementation with the *hfq* gene in pHfq/ Δhfq (Fig. 4B). These results suggest that TDH production is specifically associated with Hfq. We also examined the mRNA levels of the *tdh* gene at mid-log growth phase using quantitative real-time RT-PCR. The amount of *tdh* mRNA in the Δhfq strain was significantly higher than in the parent strain (Fig. 4C). These results suggest that the elevation of hemolytic activity in the Δhfq strain is due to activation of *tdh* gene transcription.

Transcriptional levels of the *tdh* genes in Δhfq

Previous studies showed that while two *tdh* genes (*tdhA* and *tdhS*) are located on chromosome 2 of strain RIMD2210633, in this strain TDH production was attributed only to *tdhA* [15,18]. Although we found that TDH was elevated significantly at both the transcriptional and translational levels in the Δhfq strain, quantitative real-time RT-PCR and Western blotting using anti-TDH antibody cannot distinguish between the two *tdh* genes, since their nucleotide sequences are very similar (97% identity). To solve this problem, transcriptional fusions of *tdhA* and *tdhS* to the *lacZ* reporter gene (pHN1 and pHN2, respectively; Table 1) were constructed and were monitored in Δhfq or in the isogenic parental strains. In the parental strain, the relative activity of the *tdhA-lacZ* fusion gene was much higher than the *tdhS-lacZ* fusion gene (Fig. 5). This result is consistent with previous reports [21,22]. Interestingly, the relative activity of the *tdhA-lacZ* fusion gene in the Δhfq strain was significantly elevated when compared with the parental strain, whereas there was no difference in the activity of the *tdhS-lacZ* fusion gene

Figure 2

Growth of the Δhfq strain in LB medium. Bacteria were cultured in LB medium supplemented with 3% NaCl at 37°C and the cell density was measured with a spectrophotometer. The assay was performed independently 3 times. Closed circle, parental strain; closed square, *hfq*-complemented deletion strain (pHfq Δhfq); open circle, *hfq* deletion strain (Δhfq); open square, *hfq* deletion strain with empty vector (pSN1 Δhfq).

via TTSS2, was also increased in the Δhfq strain (Fig. 6). These results indicate that the transcription of some genes encoding TTSS effector proteins is regulated in an Hfq-dependent manner.

Discussion

Hfq plays a role as a major post-transcriptional regulator in numerous bacterial pathways [9]. Previous studies have shown that Hfq affects the virulence of various pathogenic bacteria [13-18]. In this study, we have demonstrated that Hfq plays a critical role in the expression of TDH (Fig. 4). This indicates that Hfq is involved in the virulence of *V. parahaemolyticus* just as in other pathogenic bacteria, because TDH is considered to be an important virulence factor in *V. parahaemolyticus* [5-8]. The modulation of bacterial pathogenicity by Hfq has been associated with small non-coding RNAs [26-29]. It has been shown that deletion of the *hfq* gene in *P. aeruginosa* affects the expression of virulence genes by changing the stability of a small non-coding RNA [26]. Recently, Lenz *et al.* have speculated that small non-coding RNAs exist in the *V. parahaemolyticus* genome [29]. Although there is little information regarding small non-coding RNAs in *V. parahaemolyticus*, we speculate that one or more small non-coding RNAs interacting with *tdh* may regulate virulence gene expression levels.

Figure 3

Hemolytic activity in culture supernatants of parental and derivative strains. Bacteria were cultured in 10 ml of LB medium supplemented with 3% NaCl at 37°C. One hemolytic unit (HU) was defined as in a previous report [45,46]. The results are expressed as means $\pm$ SDs of 5 independent experiments. *, $p < 0.05$ versus LB; #, $p < 0.01$ versus WT. LB, negative control; WT, *V. parahaemolyticus* strain RIMD2210633 (parent strain); Δhfq , *hfq* deletion strain; pHfq/ Δhfq , *hfq*-complemented deletion strain; pSN1/ Δhfq , deletion strain with empty vector.

The production of TDH is reportedly affected by environmental factors such as pH, temperature, and chemicals [30-34], but little is known about the modulation of expression of the *tdh* gene by internal factors in *V. parahaemolyticus*. In this study, we showed that Hfq in *V. parahaemolyticus* was important for the expression of the *tdh* gene and that the transcription of the *tdhA* gene rather than the *tdhS* gene was regulated by Hfq (Figs. 4 and 5). This result indicates that the expression of the *tdh* gene is under the control of Hfq. Lin *et al.* have reported that ToxR in *V. parahaemolyticus* is associated with the production of TDH [35]. We investigated the transcriptional activity of *toxR* in the Δhfq strain under our experimental conditions, but could not find any differences between the Δhfq and wild type strains (data not shown). This suggests that the regulation of *tdh* gene expression by Hfq is *toxR*-independent. We are currently examining how Hfq regulates the expression of *tdh*.

Proteome and microarray analyses in various bacteria have shown that Hfq controls the expression of various genes [14,16,26]. Thus, Hfq is considered to be a global regulator of gene expression including genes encoding virulence factors. In this study, we confirmed that Hfq also regulates the expression of at least three effector proteins of TTSSs, namely VP1680, VopC and VopT, in *V. parahaemolyticus* (Fig. 6). It has been shown that TTSSs and

Figure 4
Production of TDH by the Δhfq strain. (A) Detection of TDH in cell lysates from mid-log phase (OD₆₀₀ = 0.6) and stationary phase (OD₆₀₀ = 1.5) cultures. (B) Detection of TDH in mid-log phase cell lysates from the parent strain and derivatives. (C) Quantification of *tdh* mRNA. Fold change in the gene transcription level from mid-log phase in the Δhfq strain versus the parent strain, as measured by quantitative real-time RT-PCR. Data analysis was as previously described [42,43]. Data are expressed as means $\pm$ SDs of 5 independent experiments. *, $p < 0.05$.

these three effector proteins contribute to pathogenicity in *V. parahaemolyticus* [23-25]. Although TTSSs in pathogenic *E. coli* and *Salmonella* are involved in the regulatory mechanisms for the expression of TTSS-related genes [36,37], this is not yet clear in *V. parahaemolyticus*. Moreover, it seems likely that other novel effector proteins will be found that contribute to the pathogenicity of *V. parahaemolyticus* [25,38]. Our results suggest that Hfq may be important for the pathogenicity of *V. parahaemolyticus*. Experiments are currently underway to analyze the mechanisms by which Hfq governs the expression of virulence genes in *V. parahaemolyticus*.

Conclusion

We have confirmed that the expression of TDH in *V. parahaemolyticus* is closely associated with the transcriptional activity of *tdhA*. Hfq regulates the transcriptional activity of *tdhA* and the expression of TDH in *V. parahaemolyticus*. We predict that Hfq may be a global regulator of pathogenicity in *V. parahaemolyticus*.

Methods

Bacteria and plasmids

The bacteria and plasmids used in this study are listed in Table 1. *V. parahaemolyticus* strain RIMD2210633 was used as the standard strain in this study [19]. *E. coli* and *V. parahaemolyticus* strains were routinely cultured in Luria-Bertani (LB) medium and LB medium supplemented with 3% NaCl, respectively.

Construction of the *hfq* deletion mutant

The mutant strain was constructed by homologous recombination [23,39]. In brief, a DNA fragment was generated by PCR using primers hfq-1 (5'-GGATCCAGCCGATAA-CATTGAGCAGGC-3') and hfq-4 (5'-CTGCAGTCAGCAC-CAGTCAAATGTAC-3') after the preparation of PCR products using primers hfq-1 and hfq-2 (5'-CTCGATGAA-GAATTAAGCCATTTATTTC-3'), and hfq-3 (5'-GAAAATAAAATGGCTTAATTCTTCATCGAG-3') and hfq-4. The amplified PCR product was cloned into the suicide vector pYAK1 [39] and the resulting plasmid was transformed into *E. coli* strain SM10 λ pir. Bacterial conjugation was performed with the recombinant *E. coli* strain SM10 λ pir and *V. parahaemolyticus*. Colonies were confirmed to have the gene deletion by PCR using the primers hfq-1 and hfq-4 and by Southern blotting.

Complementation of the *hfq* deletion mutant

Bacterial strains harboring the *hfq* gene were constructed as previously described [23]. Since the *hfq* gene promoter of *V. parahaemolyticus* has not been identified, the *tdhA* gene promoter was used for expressing *hfq* [23]. The *hfq* gene was amplified by PCR using primers hfq-5 (5'-GGATCCATGGCTAAGGGGCAATCT-3') and hfq-6 (5'-GAATTCCTTAATCTTCTGATTCTC-3'), and the PCR-generated *hfq* gene and *tdhA* promoter were cloned into pSA19CP-MCS to construct pHfq [40]. The constructed plasmid was introduced into the *V. parahaemolyticus* deletion mutant by electroporation [41].

Quantitative real-time RT-PCR

Preparation of total RNA from *V. parahaemolyticus* was carried out previously [42]. Quantitative real-time RT-PCR and data analysis were performed as previously described [42,43]. The oligonucleotide primers for *rrs*, *tdh*, *vopC* (= *vpa1321*), and *vopT* (= *vpa1327*) were as previously described [42]. The oligonucleotide primers for VP1680 were as follows: 5'-CACACTTGGTATGGCGACAG-3' and 5'-TGAGTTGGTCTTGCACTTCG-3'. The oligonucleotide primers were designed using the genome sequence of *V. parahaemolyticus* strain RIMD2210633 [19].

Preparation of polyclonal antibodies

Antibody against TDH was prepared as described previously [44]. Antibodies against VP1680, VopC, and VopT were prepared according to a previous report [23]. In

Figure 5

Activities of the *tdhA* and *tdhS* gene promoters in the WT and Δhfq strains. β -galactosidase activity measured in the parental WT strain (black bars) and the Δhfq strain (open bars). pHN1/WT, pHN1 harboring *tdhA* gene promoter in *V. parahaemolyticus* strain RIMD2210633; pHN2/WT, pHN2 harboring *tdhS* gene promoter in *V. parahaemolyticus* strain RIMD2210633; pHRP309/WT, pHRP309 in *V. parahaemolyticus* strain RIMD2210633; pHN1/ Δhfq , pHN1 harboring *tdhA* gene promoter in Δhfq strain; pHN2/ Δhfq , pHN2 harboring *tdhS* gene promoter in Δhfq strain; pHRP309/ Δhfq , pHRP309 in Δhfq strain. Data in Miller units are the means $\pm$ SDs of 5 independent experiments. *, $P < 0.01$.

brief, DNAs encoding VP1680, *vopC*, and *vopT* were generated by PCR using the following oligonucleotide primers: for VP1680, 5'-GGATCCATGGTGAATACAACGCAAAA-3' and 5'-GAATCTTAAATCCAGCCTTCGGCT-3'; for *vopC*, 5'-GGATCCATGCCAATATTAAATATTAG-3' and 5'-GAATCTTAACCATTTTCAATAGCA-3'; for *vopT*, 5'-GGATCCGTGAAGGTTTGTAGAATACA-3' and 5'-GAATCTCACTTAGCTAAATCTAGCG-3'. The resulting DNAs were cloned into pET30a. Recombinant proteins were expressed in *E. coli* BL21(DE3) and purified using His-tag affinity resins and buffer kits according to the manufacturers' instructions (Merck, Darmstadt, German). Polyclonal antibodies were generated by immunizing female New Zealand White rabbits as previously described [23].

Western blotting

V. parahaemolyticus was cultured in LB medium supplemented with 3% NaCl. Cells (2 ml of culture) were collected by centrifugation and washed once with phosphate-buffered saline (PBS; pH 7.4). The pellets were resuspended in PBS and 10 μ g of total cell lysates were used for Western blotting. After electrophoresis, proteins

were transferred to membranes and probed with each of the primary antibodies. The secondary antibody was anti-rabbit IgG conjugated to horseradish peroxidase (Bio-science, Camarillo, CA, USA). Signals were detected using an ECL Western blotting kit according to the manufacturer's instructions (GE Healthcare Bio-Science, Piscataway, NJ, USA).

Determination of hemolytic activity

Bacterial cells were prepared by diluting overnight cultures 1:1000 into fresh LB medium supplemented with 3% NaCl and incubating at 37°C for 24 h with shaking. After the elimination of bacterial cells from culture supernatants by centrifugation and a syringe filter, the supernatants were concentrated using an Amicon Ultra-15 centrifugal filter (Millipore, Billerica, MA, USA). Protein concentrations were determined using a BCA protein assay kit (Thermo Scientific, Rockford, IL, USA).

Hemolytic activity was assayed according to a previously described method [45,46] with rabbit red blood cells (rRBC) (Nippon Bio-Test Laboratories, Tokyo, Japan). Protein (20 μ g) was incubated with rRBC at 37°C for 6 h.

Table 1: Bacteria and plasmids

Bacteria and plasmids	Comments ^a	References
<i>V. parahaemolyticus</i>		
RIMD2210633	<i>tdh</i> ⁺ (serotype O3:K6; clinical isolate)	19
Δhfq	<i>hfq</i> mutant derived from RIMD2210633	This study
pHfq/ Δhfq	Δhfq harboring pHfq	This study
pSNI/ Δhfq	Δhfq harboring pSNI	This study
pHNI/WT	RIMD2210633 harboring pHNI	This study
pHN2/WT	RIMD2210633 harboring pHN2	This study
pHRP309/WT	RIMD2210633 harboring pHRP309	This study
pHNI/ Δhfq	Δhfq harboring pHNI	This study
pHN2/ Δhfq	Δhfq harboring pHN2	This study
pHRP309/ Δhfq	Δhfq harboring pHRP309	This study
<i>E. coli</i>		
SM10 λ pir	R6Kori λ pir <i>supE thi thr leu lacY supE</i>	39
Plasmids		
pYKI	Suicide vector; Cm ^r	39
pSA19CP-MCS	For gene complementation; Cm ^r	40
pHRP309	For β -galactosidase assay; <i>lacZ</i> <i>gen</i>	47
pSNI	pSA19CP-MCS containing PCR-generated <i>tdhA</i> promoter	23
pHfq	pSNI containing PCR-generated <i>hfq</i>	This study
pHNI	pHRP309 containing PCR-generated <i>tdhA</i> promoter	This study
pHN2	pHRP309 containing PCR-generated <i>tdhS</i> promoter	This study

^aCm^r, chloramphenicol resistance

Figure 6
Expression of virulence-associated genes in the Δhfq strain. (A) Western blot detection of proteins in total cell lysates. (B) Quantification of mRNAs encoding virulence-associated proteins. The fold changes in gene transcription levels in the Δhfq and parental WT strains grown to mid-log phase were determined by quantitative real-time RT-PCR. Data analysis was as previously described [42,43]. Data are the means $\pm$ SDs of 5 independent experiments. *, $p < 0.05$.

β -galactosidase assay

β -galactosidase assays were performed as described previously [47]. In brief, the *tdhA* and *tdhS* promoter regions were generated by PCR using the oligonucleotide primers *tdhA*-1 (5'-GTCGACAATTCACGACGAATCGGAG-3') and *tdhA*-2 (5'-GAATTCATATCGGTACTTCATAAA-3'), and *tdhS*-1 (5'-GTCGACAGCTTACAGCTTGGTATGCC-3') and *tdhS*-2 (5'-GAATTCGTGTTTCATAAAAAACCCAT-3'), respectively. The products were cloned into pHRP309 [48]. The constructed plasmids were introduced into *V. parahaemolyticus* strains by electroporation [41]. For this assay, bacteria were cultured in LB medium supplemented with 3% NaCl until early-log phase ($OD_{600} = 0.2$). The β -galactosidase activities of the strains were determined as described previously [49].

Data analysis

All data are expressed as means $\pm$ standard deviations. Statistical significance was calculated by paired and unpaired *t*-tests. In all cases, a *p*-value of < 0.05 was considered significant.

Authors' contributions

MN, AT, and ZS performed the experiments and drafted the manuscript. NH and KM participated in the discussions of the study design and provided technical support. YN conceived of the study and drafted the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

This work was supported by a Grant-in-Aid for the 21st Century Center of Excellent Program, Human Nutritional Science on Stress Control, The University of Tokushima.

References

- Daniels NA, MacKinnon L, Bishop R, Altekruze S, Ray B, Hammond RM, Thompson S, Wilson S, Bean NH, Griffin PM, Slutsker L: **Vibrio parahaemolyticus infections in the United States, 1973-1998.** *J Infect Dis* 2000, **181**:1661-1666.
- Blake PA, Weaver RE, Hollis DG: **Diseases of humans (other than cholera) caused by vibrios.** *Annu Rev Microbiol* 1980, **34**:341-367.
- Morris JG Jr, Black RE: **Cholera and other vibrioses in the United States.** *N Engl J Med* 1985, **312**:343-350.
- Miyamoto Y, Kato T, Obara Y, Akiyama S, Takizawa K, Yamai S: **In vitro hemolytic characteristic of Vibrio parahaemolyticus: its close correlation with human pathogenicity.** *J Bacteriol* 1969, **100**:1147-1149.
- Miyamoto Y, Obara Y, Nikkawa T, Yamai S, Kato T, Yamada Y, Ohashi M: **Simplified purification and biophysicochemical characteristic of Kanagawa phenomenon-associated hemolysin of Vibrio parahaemolyticus.** *Infect Immun* 1980, **28**:567-576.
- Naim R, Yanagihara I, Iida T, Honda T: **Vibrio parahaemolyticus thermostable direct hemolysin can induce an apoptotic cell death in Rat-1 cells from rat embryonic fibroblast cells.** *FEMS Microbiol Lett* 2001, **195**:237-244.
- Nishibuchi M, Fasano A, Russell RG, Kaper JB: **Enterotoxigenicity of Vibrio parahaemolyticus with and without encoding thermostable direct hemolysin.** *Infect Immun* 1992, **60**:3539-3545.
- Shirai H, Ito H, Hirayama T, Nakabayashi Y, Kumagai K, Takeda Y, Nishibuchi M: **Molecular epidemiologic evidence for association of thermostable direct hemolysin (TDH) and TDH-related hemolysin of Vibrio parahaemolyticus with gastroenteritis.** *Infect Immun* 1990, **58**:3568-3573.
- Valentin-Hansen P, Eriksen M, Udesen C: **The bacterial Sm-like protein Hfq: a key player in RNA transactions.** *Mol Microbiol* 2004, **51**:1525-1533.
- Vytvytska O, Jacobsen JS, Balcunaite G, Andersen JS, Baccarini M, von Gabain A: **Host factor I, Hfq, binds to Escherichia coli ompA mRNA in a growth-rate dependent fashion and regulates its stability.** *Proc Natl Acad Sci USA* 1998, **95**:14118-14123.
- Brown L, Elliot T: **Efficient translation of the RpoS sigma factor in Salmonella typhimurium requires host factor I, an RNA-binding protein encoded by the hfq gene.** *J Bacteriol* 1996, **178**:3763-3770.
- Zhang A, Altuvia S, Tiwari A, Argaman L, Hegge-Aronis R, Storz G: **The OxyS regulatory RNA represses RpoS translation and binds the Hfq (HF-I) protein.** *EMBO J* 1998, **17**:6061-6068.
- Christiansen JK, Larsen MH, Ingmer H, Sogaard-Andersen L, Kallipolitis BH: **The RNA-binding protein Hfq of Listeria monocytogenes: role in stress tolerance and virulence.** *J Bacteriol* 2004, **186**:3355-3362.
- Ding Y, Davis BM, Waldor MK: **Hfq is essential for Vibrio cholerae virulence and down regulates sigma expression.** *Mol Microbiol* 2004, **53**:345-354.
- Sonnleitner E, Hagens S, Rosenau F, Wilhelm S, Habel A, Jager KE, Bläsi U: **Reduced virulence of a hfq mutant of Pseudomonas aeruginosa O1.** *Microb Pathog* 2003, **35**:217-228.
- Sittka A, Pfeiffer V, Tedin K, Vogel J: **The RNA chaperone Hfq is essential for the virulence of Salmonella typhimurium.** *Mol Microbiol* 2007, **63**:193-217.
- Nakao H, Watanabe H, Nakayama S, Tekada T: **yst gene expression in Yersinia enterocolitica is positively regulated by a chromosomal region that is highly homologous to Escherichia coli host factor I gene (hfq).** *Mol Microbiol* 1995, **18**:859-865.
- Robertson GT, Roop RM Jr: **The Brucella abortus host factor I (HF-I) protein contributed to stress resistance during stationary phase and is a major determinant of virulence in mice.** *Mol Microbiol* 1999, **34**:690-700.
- Makino K, Oshima K, Kurokawa K, Yokoyama K, Uda T, Tagomori K, Iijima Y, Najima M, Nakano M, Yamashita A, Kubota Y, Kimura S, Yasunaga T, Honda T, Shinagawa H, Hattori M, Iida T: **Genome sequence of Vibrio parahaemolyticus: a pathogenic mechanism distinct from that of V. cholerae.** *Lancet* 2002, **361**:743-749.
- Tsui HC, Leung HC, Winkler ME: **Characterization of broadly pleiotropic phenotypes caused by an hfq insertion mutation in Escherichia coli K-12.** *Mol Microbiol* 1994, **13**:35-49.
- Park KS, Ono T, Rokuda M, Jang MH, Iida T, Honda T: **Cytotoxicity and enterotoxigenicity of the thermostable direct hemolysin-deletion mutants of Vibrio parahaemolyticus.** *Microbiol Immunol* 2004, **48**:313-318.
- Okuda J, Nishibuchi M: **Manifestation of the Kanagawa phenomenon, the virulence-associated phenotype, of Vibrio parahaemolyticus depends on a particular single base change in the promoter of the thermostable direct haemolysin gene.** *Mol Microbiol* 1998, **30**:499-511.
- Park KS, Ono T, Rokuda M, Jang MH, Okada K, Iida T, Honda T: **Functional characterization of type III secretion systems of Vibrio parahaemolyticus.** *Infect Immun* 2004, **72**:6659-6665.
- Kodama T, Rokuda M, Park KS, Cantarelli VV, Matsuda S, Iida T, Honda T: **Identification and characterization of VopT, a novel ADP-ribosyltransferase effector protein secreted via the Vibrio parahaemolyticus type III secretion system 2.** *Cell Microbiol* 2007, **9**:2598-2609.
- Ono T, Park KS, Ueta M, Iida T, Honda T: **Identification of proteins secreted via Vibrio parahaemolyticus type III secretion system 1.** *Infect Immun* 2006, **74**:1032-1042.
- Sonnleitner E, Schuster M, Sorger-Domenigg T, Greenberg EP, Bläsi U: **Hfq-dependent alterations of the transcriptome profile and effects on quorum sensing in Pseudomonas aeruginosa.** *Mol Microbiol* 2006, **59**:1542-1558.
- Pfeiffer V, Sittka A, Tomer R, Tedin K, Brinkmann V, Vogel J: **A small non-coding RNA of the invasion gene island (SPI-1) represses outer membrane protein synthesis from the Salmonella core genome.** *Mol Microbiol* 2007, **66**:1174-1191.
- Kay E, Humair B, Denervaud V, Riedel K, Spahr S, Eberl L, Valverde C, Haas D: **Two GacA-dependent small RNAs modulate the quorum-sensing response in Pseudomonas aeruginosa.** *J Bacteriol* 2006, **188**:6026-6033.
- Lenz DH, Mok KC, Lilley BN, Kulkarni RV, Wingreen NS, Bassler BL: **The small RNA chaperone Hfq and multiple small RNAs control quorum sensing in Vibrio harvey and Vibrio cholerae.** *Cell* 2004, **118**:69-82.
- Cherwonogrodzky JW, Clark AG: **Effect of pH on the production of the Kanagawa hemolysin by Vibrio parahaemolyticus.** *Infect Immun* 1981, **34**:115-119.
- Karunasagar I: **Production of hemolysin by Vibrio parahaemolyticus in a chemically defined medium.** *Appl Environ Microbiol* 1981, **41**:1272-1275.
- Cherwonogrodzky JW, Skinner MA, Clark G: **Effect of D-tryptophan on hemolysin production in Vibrio parahaemolyticus.** *J Clin Microbiol* 1984, **20**:909-911.
- Wong HC, Lee YS: **Regulation of iron on bacterial growth and production of thermostable direct hemolysin by Vibrio parahaemolyticus in intraperitoneal infected mice.** *Microbiol Immunol* 1994, **38**:367-371.
- Osawa R, Yamai S: **Production of thermostable direct hemolysin by Vibrio parahaemolyticus enhanced by conjugated bile acids.** *Appl Environ Microbiol* 1996, **62**:3023-3025.
- Lin Z, Kumagai K, Baba K, Mekalanos JJ, Nishibuchi M: **Vibrio parahaemolyticus has a homolog of the Vibrio cholerae toxRS operon that mediates environmentally induced regulation of the thermostable direct hemolysin gene.** *J Bacteriol* 1993, **175**:3844-3855.
- Ellermeier JR, Schlauch JM: **Adaptation to the host environment: regulation of the SPII type III secretion system in Salmonella enterica serovar Typhimurium.** *Curr Opin Microbiol* 2007, **10**:24-29.
- Mellies JL, Elliott SJ, Sperandio V, Donnenberg MS, Kaper JB: **The Per regulon of enteropathogenic Escherichia coli: identification of a regulatory cascade and a novel transcriptional activator, the locus of enterocyte effacement (LEE)-encoded regulator (Ler).** *Mol Microbiol* 1999, **33**:296-306.
- Liverman ADB, Cheng HC, Trosky JE, Leung DW, Yarbrough ML, Burdette DL, Rosen MK, Orth K: **Arp 2/3-independent assembly of actin by Vibrio type III effector VopL.** *Proc Natl Acad Sci USA* 2007, **104**:17117-17122.
- Kodama T, Akeida Y, Kono G, Takahashi A, Imura K, Iida T, Honda T: **The EspB protein of enterohaemorrhagic Escherichia coli interacts directly with α -catenin.** *Cell Microbiol* 2002, **4**:213-222.
- Nomura T, Hamashima H, Okamoto K: **Carboxy terminal region of haemolysin of Aeromonas sobria triggers dimerization.** *Microbiol Pathog* 2000, **28**:25-36.
- Hamashima H, Nakano T, Tamura S, Arai T: **Genetic transformation of Vibrio parahaemolyticus, Vibrio alginolyticus, and Vibrio**

- cholerae* non O-1 with plasmid DNA by electroporation. *Microbiol Immunol* 1990, **34**:703-708.
42. Nakano M, Takahashi A, Sakai Y, Nakaya Y: **Modulation of pathogenicity with norepinephrine related to the type III secretion system of *Vibrio parahaemolyticus*.** *J Infect Dis* 2007, **195**:1353-1360.
 43. Leverton LQ, Kaper JB: **Temporal expression of enteropathogenic *Escherichia coli* virulence genes in an in vitro model of infection.** *Infect Immun* 2005, **73**:1034-1043.
 44. Takahashi A, Yamamoto C, Kodama T, Yamashita K, Harada N, Nakano M, Honda T, Nakaya Y: **Pore formation of thermostable direct hemolysin secreted from *Vibrio parahaemolyticus* in lipid bilayers.** *Int J Toxicol* 2006, **25**:409-418.
 45. Tang GQ, Iida T, Yamamoto K, Honda T: **A mutant toxin of *Vibrio parahaemolyticus* thermostable direct hemolysin which has lost hemolytic activity but retains ability to bind to erythrocytes.** *Infect Immun* 1994, **62**:3299-3304.
 46. Honda T, Ni YX, Miwatani T: **Purification and characterization of a hemolysin produced by a clinical isolate of Kanagawa phenomenon-negative *Vibrio parahaemolyticus* and related to the thermostable direct hemolysin.** *Infect Immun* 1988, **56**:961-965.
 47. Shime-Hattori A, Iida T, Arita M, Park KS, Kodama T, Honda T: **Two type IV pili of *Vibrio parahaemolyticus* play different roles in biofilm formation.** *FEMS Microbiol Lett* 2006, **264**:89-97.
 48. Parales RE, Harwood CS: **Construction and use of a new broad-host range *lacZ* transcriptional fusion vector, pHRP309, for Gram⁻ bacteria.** *Gene* 1993, **133**:23-30.
 49. Miller JH: **Experiment 48 Assay of β -Galactosidase.** In *Experiments in Molecular Genetics* Edited by: Miller J. New York: Cold Spring Harbor Laboratory; 1972:352-355.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp