

RESEARCH ARTICLE

Open Access

A comparison of conventional methods for the quantification of bacterial cells after exposure to metal oxide nanoparticles

Hongmiao Pan¹, Yongbin Zhang², Gui-Xin He³, Namrata Katagori³ and Huizhong Chen^{1*}

Abstract

Background: Due to potential interference of nanoparticles on bacterial quantification, there is a challenge to develop a fast, accurate and reproducible method for bacterial quantification. Currently various bacterial quantification methods are used by researchers performing nanoparticles study, but there has been no efficacy evaluation of these methods. Here we study interference of nanoparticles on three most commonly used conventional bacterial quantification methods, including colony counting to determine the colony-forming units (CFU), spectrophotometer method of optical density (OD) measurement, and flow cytometry (FCM).

Results: Three oxide nanoparticles including ZnO, TiO₂, and SiO₂ and four bacterial species including *Salmonella enterica* serovar Newport, *Staphylococcus epidermidis*, *Enterococcus faecalis*, and *Escherichia coli* were included in the test. Results showed that there is no apparent interference of the oxide nanoparticles on quantifications of all four bacterial species by FCM measurement; CFU counting is time consuming, less accurate and not suitable for automation; and the spectrophotometer method using OD measurement was the most unreliable method to quantify and detect the bacteria in the presence of the nanoparticles.

Conclusion: In summary, FCM measurement proved to be the best method, which is suitable for rapid, accurate and automatic detection of bacteria in the presence of the nanoparticles.

Keywords: Nanoparticles, Bacterial quantification, Flow cytometry (FCM), Colony-forming units (CFU), Optical density (OD)

Background

Nanoparticles (NPs) offer spectacular properties to their bulk materials, such as a high surface area to volume ratio, new mechanical, chemical, electrical, optical, magnetic, electro-optical, and magneto-optical properties [1]. Nanotechnology is one of the fastest growing areas of the high tech economy [2,3]. Products using nanoparticles - also known as nanomaterials (particle sizes less than 100 nm)-can be found in almost every area of our daily lives, from cosmetics to clothing to foods to drug products [4-7]. There are hundreds of cosmetics that contain nanomaterials, such as ZnO, TiO₂, and SiO₂, in

the market now and the number of these products are increasing rapidly [8]. Nanoscale materials can find use in many areas related to the food industry including agriculture, food processing, food security, packaging, nutrition and nutraceuticals [9-11]. Nanoscale materials have been used as novel antimicrobial agents [12]. Due to their powerful antimicrobial activity and particular modes of action, nanoparticles provide an attractive alternative to classic antibiotics in the development of next-generation antibiotic agents [13-15]. Various antimicrobial nanoparticles and nanosized carriers for antibiotics delivery have been developed to effectively treat infectious diseases, especially those caused by antibiotic-resistant microorganisms [16,17]. Nanoparticles behave differently from their respective bulk materials and thus

* Correspondence: huizhong.chen@fda.hhs.gov

¹Division of Microbiology, National Center for Toxicological Research, U.S.

Food and Drug Administration, Jefferson, AR 72079, USA

Full list of author information is available at the end of the article

the unique properties of the nanoparticles might also cause adverse health effects on human, animal and environment. The speedy commercialization of nanotechnology requires thoughtful and careful environmental, animal and human health safety assessment [18,19].

The detection and quantification of viable bacteria plays a critical role in quality control programs of the food, cosmetics and drug industry to prevent illness and in clinical diagnosis and therapeutics. Currently there are many methods used for the detection and quantification of bacteria, including conventional and molecular approaches [20-24]. Conventionally identification of bacteria is usually performed by three methods including culture-based counting for colony-forming units (CFU) [22,25], spectrophotometer method of optical density (OD) measurement [23,24], and flow cytometry (FCM) [26,27]. In spite of the sensitivity and reliability, counting CFU is time-consuming and labor-intensive [28,29]. CFU determination is the conventional method to quantify bacteria, but only detects those that are able to grow on specific solid media, which excludes the detection of unculturable live, inactive or damaged bacterial cells [30,31]. Therefore, CFU counting tends to undercount the actual number of the bacteria. For example, anaerobic bacteria are not able to grow on the media and cultural conditions suitable for growth of aerobic bacteria. Optical density method measures turbidity associated directly with bacterial growth which is rapid, low cost and non-destructive, however, it measures live as well as dead bacterial cell debris. Flow cytometry is a relatively newly developed technique and enables a fast and reliable detection of all bacteria including the non-cultivable microorganisms. It enables researchers to reliably distinguish and quantitate live and dead bacteria with the aid of a flow cytometer in a mixed population containing various bacterial types [32]. Besides, Flow cytometry method is able to provide morphometric and functional properties of the detected bacteria [33,34]. Currently all these three methods are employed to quantify bacterial contents in the presence of nanoparticles [35-39]. So far there has not been any research performed concerning potential interference by nanoparticles on the bacterial counting methods.

The aim of this study was to compare three commonly used conventional methods for bacterial detection and quantification in the presence of nanoparticles. In this study, three nanoparticles, ZnO, TiO₂, and SiO₂, which are commonly used in commercial products and four important human pathogens (*Salmonella enterica* serovar Newport, *Staphylococcus epidermidis*, *Enterococcus faecalis*, and *Escherichia coli*) frequently found in various products and environment representing Gram-positive and Gram-negative bacteria were employed for this study.

Results and discussion

Physical and chemical characterizations of nanomaterials

It is critical to conduct physical and chemical characterization of testing nanomaterials in nanotechnology research. Size, size distribution, surface charge, aggregation or agglomeration status, and shape have been considered as the most important parameters for nanomaterials. We evaluated these parameters using TEM and Zetasizer as described in the material and methods section. TEM analysis indicated that the ZnO, TiO₂ and SiO₂ nanoparticles have spherical shape with slightly agglomeration (Figure 1). The primary size of ZnO, TiO₂ and SiO₂ nanoparticles were measured as 14.0 ± 4.9 nm, 19.7 ± 5.7 nm and 17.4 ± 5.1 nm, respectively (Table 1). The range of the diameter of the ZnO, TiO₂ and SiO₂ nanoparticle was 6.3-30.5 nm, 10.2-31.2 nm and 8.0-27.9 nm. Zetasizer analysis indicated that the average size of ZnO, TiO₂ and SiO₂ nanoparticles in buffer solution was 2308.3 ± 159.1 nm, 2738.3 ± 303.3 nm and 915.0 ± 35.8 nm (mean \pm SD). The average surface charge of the ZnO, TiO₂, SiO₂ nanoparticles in buffer solution was 17.6 ± 0.7 mV, 27.2 ± 3.1 mV, -5.7 ± 0.4 mV, respectively (Table 1). TEM directly measured the primary size of the nanoparticles based on the projected area; while Dynamic Light Scattering (DLS) measured the hydrodynamic diameter of the nanoparticles based on the translational diffusion area of the particle being measured. The same samples of these nanoparticles in buffer were measured with a bigger size by zetasizer analysis than the measurement using TEM. This is due to the differences in the weighted averages determined by these two techniques, and also the differences in the physical properties measured. TEM is sensitive to the size of primary particles, whereas DLS is sensitive to the presence of small quantities of large particles or aggregates.

Effect of concentrations of nanoparticle on quantification of bacteria

Potential interference of nanoparticles on the quantification of bacteria poses a challenge for the detection of bacterial contaminations in various consumer products. Therefore, development of a rapid, sensitive and accurate method for detection of bacteria in the presence of nanoparticles is crucial for food, drug, cosmetic and other consumable products. Among many bacterial identification and quantification methods, three of them including culture-based counting for CFU, spectrophotometer method of optical density measurement, and more recently flow cytometry are commonly used. ZnO, TiO₂, and SiO₂ have been found in many commercial products including food, food supplements, cosmetics and drugs. *S. enterica* Newport, *S. epidermidis*, *E. faecalis*, and *E. coli*, which are important human pathogens, are good representatives for Gram-positive and Gram-negative bacteria (Table 2). In this experiment the effect of




Figure 1 Characterization of ZnO, TiO₂, or SiO₂ nanoparticles by transmission electron microscopy (TEM). Nanoparticles were deposited on formvar carbon coated grids and dried for TEM imaging. Images were analyzed in high resolution mode with an acceleration voltage of 100 kV. Morphology of ZnO, TiO₂ or SiO₂ is shown in left, middle and right of the above images. Scale Bar = 20 nm.

various concentrations of nanoparticles on quantification of *S. enterica* Newport, *S. epidermidis*, *E. faecalis*, and *E. coli* was investigated by exposing 5 ml of samples containing approximately 10⁹ cells/ml to various concentrations of ZnO, TiO₂, and SiO₂ (0, 0.1, 0.2, 0.3, 0.5, and 1 mg/ml final concentration) for 1 hr, respectively (Table 3). As shown in Table 3, with increasing concentrations of ZnO, TiO₂, and SiO₂, there was no apparent interference of the nanoparticles on quantifications of all four bacterial species by flow cytometry measurement using the BacLight LIVE/DEAD bacterial viability and counting kit. As shown in Figure 2 as example, two distinctive groups were formed. Group P2 was the population of living bacterial cells, while group P3 was the population of dead bacterial cells at the presence of 0.2 mg/ml nanoparticles. Compared to a control, which did not contain nanoparticles, no shifts of the bacterial population or background increase were observed (Figure 2). Since more than 20,000 bacterial cells per sample were counted by flow cytometry measurement, high accuracy and excellent reproducibility of the quantification was achieved for both live and dead bacterial cells (Table 3). Although no apparent interference of the nanoparticles on quantifications of all four bacterial species was observed by using CFU counting, it was a time consuming and labor intensive procedure. Besides, it took long time training and practice for mastering the technique of dilution in order to get reliable counts from one batch to another and from one plate to another in CFU counting. Furthermore, the data obtained by CFU measurement is less accurate and reproducible due to a limited number of bacterial cells counted (several hundred bacterial colonies counted (Table 3). The decreasing numbers of

Table 1 Characterization of TiO₂, ZnO, and SiO₂ nanoparticles in Milli-Q water solutions

Physical Parameters	ZnO	TiO ₂	SiO ₂
Primary size (nm)	14.0 ± 4.9	19.7 ± 5.7	17.4 ± 5.1
Primary size range(nm)	6.3 – 30.5	10.2 - 31.2	8.0 – 27.9
Hydrodynamic size (nm)	2738.3 ± 303.3	2308.3 ± 159.1	915.0 ± 35.8
Shape	spherical	spherical	spherical
Agglomerate in solution	Yes	Yes	Yes
Zeta potential ζ (mV)	17.6 ± 0.7	27.2 ± 3.1	-5.7 ± 0.4

Table 2 Bacterial species used in the study

Species name	Gram ¹	Culture condition	Isolation
<i>Salmonella enterica</i> serovar Newport	-	aerobic	human intestine
<i>Staphylococcus epidermidis</i> ATCC 12228	+	aerobic	human skin
<i>Enterococcus faecalis</i> ATCC 27274	+	anaerobic	human intestine
<i>Escherichia coli</i> ATCC 25922	-	anaerobic	human intestine

¹+, Gram-positive; -, Gram-negative.

the bacteria by using CFU and flow cytometry were resulted from antibacterial effects caused by both nanoparticles TiO₂ and ZnO. As shown in Table 3, nanoparticles had adverse effect on quantification of bacteria using the spectrophotometer method of optical density measurement with severity of TiO₂ > ZnO > SiO₂. For example, in the presence of 0.1 mg TiO₂, the number of *S. epidermidis* cells was not detectable due to high background interference from the nanoparticles in the samples. With 0.3 mg/ml TiO₂, *S. enterica* Newport and *E. faecalis* cells were 45.2 and 42.8% of those measured by FCA, respectively. On the other hand, *E. coli* cells were more than 8-folds than that by FCA in the presence of 0.3 mg/ml TiO₂.

Effect of bacterial concentrations on quantification of bacteria after exposure to nanoparticles

In this experiment, we further investigated interference of nanoparticles ZnO (0.5 mg/ml), TiO₂ (0.5 mg/ml), and SiO₂ (1 mg/ml) on quantifications of *S. enterica* Newport, *S. epidermidis*, *E. faecalis*, and *E. coli* with various bacterial concentrations by comparison of flow cytometry with optical density measurement. As shown in Table 4, with the decreasing concentrations of the tested bacteria, no apparent interference of the nanoparticles on quantifications of all four bacterial species by flow cytometry was observed. However, with decreasing concentrations of the tested bacteria, the adverse effect on quantification of bacteria was dramatically increased using the spectrophotometer method of optical density measurement, which reflected a progressively worse estimate of the bacterial counts as the ratio of numbers of bacteria and nanoparticles in the suspension decreased. For example, in the presence of 0.1 mg TiO₂, number of *S. enterica* Newport cells could not be detected due to high background interference from the nanoparticles in the samples. *S. enterica* Newport, *S. epidermidis*, *E. faecalis* could not be quantified in the presence of 0.5 mg/ml TiO₂. The data obtained from the bacterial quantification in the presence of 0.5 mg/ml of ZnO were either not able to be detected or not accurate. Due to lower interference of SiO₂ at 1 mg/ml on the bacterial quantification, there was no apparent

difference between flow cytometry and optical density measurement (Table 4).

Conclusions

In summary, this study compared three most commonly used bacterial quantification methods including colony counts, spectrophotometer method of optical density measurement, and flow cytometry in the presence of metal oxide nanoparticles. Our results demonstrated that flow cytometry is the best method with no apparent interference by the nanoparticles, indicating that it is suitable for rapid, accurate and automatic detection of bacteria. Flow cytometry is also able to detect both live and dead bacterial cells and allows detection of all bacteria including those that are uncultured. Although the bacterial quantification determined by plate counts was not affected by the nanoparticles, it was time consuming, less accurate and not suitable for automation. The spectrophotometer method using optical density measurement was the most unreliable method to quantify and detect bacteria in the presence of oxide nanoparticles. The data presented in this study indicated that flow cytometry method for bacterial quantification is superior to the other two methods. This study provides data examining the potential interference of oxide nanoparticles on bacterial quantification. The information provided here will be useful in the assessment of bacterial contamination in food, drug and cosmetic products containing nanoparticles. Future studies on other nanoparticles and limit of the bacterial detection by FMC are warranted.

Methods

Materials and preparation of nanoparticle suspensions

ZnO (purity >97%), TiO₂ (purity ≥99.5%), and SiO₂ (purity 99.5%) nanoparticles were purchased from Sigma Chemical Co. The LIVE/DEAD BacLight bacterial viability and counting kit containing solutions of 3.34 mM SYTO9 in dimethyl sulfoxide (DMSO, 200 μl), 20 mM propidium iodide (PI) in DMSO (200 μl) and a calibrated suspension of microspheres (diameter: 6 μm, 1 ml; concentration: 1.0 × 10⁸ beads/ml) and SYTO 9 green fluorescent nucleic acid stain (5 mM solution in DMSO, 100 μl) were purchased from Molecular Probes (Eugene, Oregon). Suspensions of the nanoparticles were prepared with Milli-Q water by means of ultrasonic vibration in a BRANSON 3200 UltraSonic Cleaner for 30 min and the stock solutions were vortexed briefly before each use [40-42].

Physical and chemical characterizations of nanomaterials

The size, shape and morphology of ZnO, TiO₂ or SiO₂ nanoparticles were determined using transmission electron microscopy (TEM). The nanoparticles were homogeneously

Table 3 Interference of oxide nanoparticles on bacterial quantification

Nanoparticles (mg/ml)	ZnO				TiO ₂				SiO ₂			
	<i>S. enterica</i> Newport (cells/ml) ^a				<i>S. enterica</i> Newport (cells/ml)				<i>S. enterica</i> Newport (cells/ml)			
	FMC		CFU	OD ₆₆₀ ^b	FMC		CFU	OD ₆₆₀	FMC		CFU	OD ₆₆₀
	Total	Live			Total	Live			Total	Live		
0	1.37 × 10 ⁹	1.36 × 10 ⁹	8.17 × 10 ⁸	1.37 × 10 ⁹	1.23 × 10 ⁹	1.22 × 10 ⁹	1.18 × 10 ⁹	1.23 × 10 ⁹	1.28 × 10 ⁹	1.26 × 10 ⁹	6.32 × 10 ⁸	1.28 × 10 ⁹
0.1	1.31 × 10 ⁹	1.30 × 10 ⁹	1.00 × 10 ⁹	1.46 × 10 ⁹	1.00 × 10 ⁹	9.94 × 10 ⁸	7.00 × 10 ⁸	9.16 × 10 ⁸	1.23 × 10 ⁹	1.22 × 10 ⁹	6.50 × 10 ⁸	1.20 × 10 ⁹
0.2	1.29 × 10 ⁹	1.28 × 10 ⁹	5.83 × 10 ⁸	1.28 × 10 ⁹	8.15 × 10 ⁸	8.05 × 10 ⁸	5.67 × 10 ⁸	5.89 × 10 ⁸	1.22 × 10 ⁹	1.20 × 10 ⁹	5.83 × 10 ⁸	1.18 × 10 ⁹
0.3	1.27 × 10 ⁹	1.14 × 10 ⁹	7.00 × 10 ⁸	1.19 × 10 ⁹	7.14 × 10 ⁸	7.06 × 10 ⁸	5.50 × 10 ⁸	3.23 × 10 ⁸	1.20 × 10 ⁹	1.18 × 10 ⁹	5.83 × 10 ⁸	1.16 × 10 ⁹
0.5	1.23 × 10 ⁹	1.21 × 10 ⁹	6.33 × 10 ⁸	1.01 × 10 ⁹	4.26 × 10 ⁸	4.13 × 10 ⁸	4.33 × 10 ⁸	- ^c	1.24 × 10 ⁹	1.21 × 10 ⁹	5.67 × 10 ⁸	1.15 × 10 ⁹
1	1.12 × 10 ⁹	1.10 × 10 ⁹	5.00 × 10 ⁸	7.15 × 10 ⁸	2.41 × 10 ⁸	2.35 × 10 ⁸	1.50 × 10 ⁸	-	1.22 × 10 ⁹	1.20 × 10 ⁹	7.17 × 10 ⁸	1.09 × 10 ⁹
	<i>S. epidermidis</i> ATCC 12228 (cells/ml)				<i>S. epidermidis</i> ATCC 12228 (cells/ml)				<i>S. epidermidis</i> ATCC 12228 (cells/ml)			
0	3.53 × 10 ⁸	3.46 × 10 ⁸	9.33 × 10 ⁷	3.53 × 10 ⁸	4.46 × 10 ⁸	4.40 × 10 ⁸	1.20 × 10 ⁸	4.46 × 10 ⁸	5.20 × 10 ⁸	4.74 × 10 ⁸	2.00 × 10 ⁸	5.20 × 10 ⁸
0.1	2.13 × 10 ⁸	1.94 × 10 ⁸	2.18 × 10 ⁷	2.73 × 10 ⁸	1.21 × 10 ⁸	1.19 × 10 ⁸	2.00 × 10 ⁷	-	1.06 × 10 ⁸	9.57 × 10 ⁷	1.18 × 10 ⁸	4.48 × 10 ⁸
0.2	1.37 × 10 ⁸	1.18 × 10 ⁸	1.63 × 10 ⁷	1.23 × 10 ⁸	2.65 × 10 ⁷	2.62 × 10 ⁷	2.00 × 10 ⁷	-	7.27 × 10 ⁷	6.55 × 10 ⁷	6.50 × 10 ⁷	4.54 × 10 ⁸
0.3	1.71 × 10 ⁷	1.45 × 10 ⁷	1.37 × 10 ⁷	3.20 × 10 ⁸	1.46 × 10 ⁷	1.44 × 10 ⁷	3.33 × 10 ⁷	-	5.13 × 10 ⁷	4.60 × 10 ⁷	5.00 × 10 ⁷	5.00 × 10 ⁸
0.5	1.65 × 10 ⁷	1.45 × 10 ⁷	1.33 × 10 ⁷	1.85 × 10 ⁸	6.47 × 10 ⁶	6.40 × 10 ⁶	5.83 × 10 ⁷	-	6.72 × 10 ⁷	6.32 × 10 ⁷	5.83 × 10 ⁷	4.75 × 10 ⁸
1	3.31 × 10 ⁷	3.00 × 10 ⁷	1.10 × 10 ⁷	-	6.20 × 10 ⁷	6.11 × 10 ⁷	1.07 × 10 ⁸	-	2.21 × 10 ⁸	2.04 × 10 ⁸	1.18 × 10 ⁸	4.84 × 10 ⁸
	<i>E. faecalis</i> ATCC 27274 (cells/ml)				<i>E. faecalis</i> ATCC 27274 (cells/ml)				<i>E. faecalis</i> ATCC 27274 (cells/ml)			
0	2.29 × 10 ⁹	2.28 × 10 ⁹	1.17 × 10 ⁹	2.29 × 10 ⁹	2.21 × 10 ⁹	2.17 × 10 ⁹	1.07 × 10 ⁹	2.21 × 10 ⁹	2.47 × 10 ⁹	2.42 × 10 ⁹	1.87 × 10 ⁹	2.47 × 10 ⁹
0.1	2.14 × 10 ⁹	2.06 × 10 ⁹	1.50 × 10 ⁹	2.13 × 10 ⁹	1.84 × 10 ⁹	1.81 × 10 ⁹	9.50 × 10 ⁸	1.69 × 10 ⁹	2.17 × 10 ⁹	2.11 × 10 ⁹	1.22 × 10 ⁹	2.40 × 10 ⁹
0.2	2.0 × 10 ⁹	1.92 × 10 ⁹	1.42 × 10 ⁹	1.73 × 10 ⁹	1.42 × 10 ⁹	1.40 × 10 ⁹	5.50 × 10 ⁸	1.07 × 10 ⁹	1.64 × 10 ⁹	1.61 × 10 ⁹	1.18 × 10 ⁹	2.30 × 10 ⁹
0.3	1.47 × 10 ⁹	1.44 × 10 ⁹	1.28 × 10 ⁹	1.54 × 10 ⁹	1.23 × 10 ⁹	1.22 × 10 ⁹	3.80 × 10 ⁸	5.26 × 10 ⁸	1.34 × 10 ⁹	1.33 × 10 ⁹	1.14 × 10 ⁹	2.23 × 10 ⁹
0.5	1.45 × 10 ⁹	1.40 × 10 ⁹	1.15 × 10 ⁹	8.57 × 10 ⁸	5.58 × 10 ⁸	5.54 × 10 ⁸	1.30 × 10 ⁸	-	8.69 × 10 ⁸	8.59 × 10 ⁸	7.00 × 10 ⁸	2.10 × 10 ⁹
1	1.07 × 10 ⁹	1.03 × 10 ⁹	7.00 × 10 ⁸	-	1.70 × 10 ⁶	1.60 × 10 ⁶	2.95 × 10 ⁶	-	4.44 × 10 ⁸	4.33 × 10 ⁸	5.00 × 10 ⁸	1.90 × 10 ⁹
	<i>E. coli</i> ATCC 25922 (cells/ml)				<i>E. coli</i> ATCC 25922 (cells/ml)				<i>E. coli</i> ATCC 25922 (cells/ml)			
0	6.56 × 10 ⁸	5.64 × 10 ⁸	3.98 × 10 ⁸	6.65 × 10 ⁸	6.41 × 10 ⁸	6.32 × 10 ⁸	6.83 × 10 ⁸	6.41 × 10 ⁸	5.52 × 10 ⁸	5.46 × 10 ⁸	5.67 × 10 ⁸	5.52 × 10 ⁸
0.1	5.22 × 10 ⁸	4.95 × 10 ⁸	3.93 × 10 ⁸	6.18 × 10 ⁸	3.28 × 10 ⁸	3.26 × 10 ⁸	8.33 × 10 ⁷	4.86 × 10 ⁸	3.73 × 10 ⁸	3.68 × 10 ⁸	2.83 × 10 ⁸	5.21 × 10 ⁸
0.2	4.50 × 10 ⁸	4.17 × 10 ⁸	3.88 × 10 ⁸	5.56 × 10 ⁸	7.67 × 10 ⁷	7.61 × 10 ⁷	1.17 × 10 ⁷	3.07 × 10 ⁸	2.52 × 10 ⁸	2.49 × 10 ⁸	2.17 × 10 ⁸	5.08 × 10 ⁸
0.3	3.65 × 10 ⁸	3.54 × 10 ⁸	3.87 × 10 ⁸	4.97 × 10 ⁸	1.90 × 10 ⁷	1.88 × 10 ⁷	1.17 × 10 ⁷	1.63 × 10 ⁸	2.19 × 10 ⁸	2.16 × 10 ⁸	1.50 × 10 ⁸	5.11 × 10 ⁸
0.5	1.36 × 10 ⁸	1.17 × 10 ⁸	2.93 × 10 ⁸	2.89 × 10 ⁸	7.13 × 10 ⁶	6.97 × 10 ⁶	9.02 × 10 ⁶	-	2.03 × 10 ⁸	2.02 × 10 ⁸	2.50 × 10 ⁸	4.76 × 10 ⁸
1	1.43 × 10 ⁸	1.37 × 10 ⁸	3.10 × 10 ⁸	1.59 × 10 ⁸	2.21 × 10 ⁷	2.18 × 10 ⁷	4.58 × 10 ⁷	-	2.38 × 10 ⁸	2.37 × 10 ⁸	2.83 × 10 ⁸	4.67 × 10 ⁸

^aBacterial cell concentrations were measured by flow cytometry (FCM), culture-based counting for colony-forming units (CFU), and spectrophotometer method of optical density (OD) measurement after 1 hr exposure to different concentrations of ZnO, TiO₂ and SiO₂ nanoparticles; inoculum used for each experiment was indicated in the control samples, i.e. no nanoparticles.

^bPresented data were converted from each sample cell concentration according to the each species standard curve of cell/ml vs OD₆₆₀ and as mean of triplicate with standard deviations (SD) of < 5% from FCM and OD₆₆₀ and <10% from CFU. Inoculum used for each experiment was indicated in the control samples, i.e. no nanoparticles.

^cValue was negative.




Figure 2 Examples of flow cytometric for *E. faecalis* exposure to nanoparticles-ZnO, TiO₂, and SiO₂ at concentration of 0.2 mg/ml. Fluorescence (FL1-H/FL3-H) was tested from bacterial cells inside gate P1 in a FSC-H (forward scatter-H)/SSC-H (side scatter-H) density plots. Live bacterial cells (gate P2); dead bacterial cells (gate P3).

Table 4 Quantification of bacterial cells at various concentrations in the presence of oxide nanoparticles

Strain name	Control (No nanoparticles) ^a			ZnO (0.5 mg/ml)			TiO ₂ (0.5 mg/ml)			SiO ₂ (1 mg/ml)		
	FCM		OD ₆₆₀ ^b	FCM		OD ₆₆₀	FCM		OD ₆₆₀	FCM		OD ₆₆₀
	Total cell no.	Live cell no.		Total cell no.	Live cell no.		Total cell no.	Live cell no.		Total cell no.	Live cell no.	
<i>S. enterica</i> Newport	1.34 × 10 ⁹	1.31 × 10 ⁹	1.34 × 10 ⁹	1.20 × 10 ⁹	1.17 × 10 ⁹	7.47 × 10 ⁸	4.72 × 10 ⁸	4.63 × 10 ⁸	-	1.29 × 10 ⁹	1.29 × 10 ⁹	1.36 × 10 ⁹
	6.76 × 10 ⁸	6.61 × 10 ⁸	7.45 × 10 ⁸	5.18 × 10 ⁸	5.06 × 10 ⁸	1.73 × 10 ⁸	9.58 × 10 ⁷	9.21 × 10 ⁷	-	6.07 × 10 ⁸	6.06 × 10 ⁸	7.91 × 10 ⁷
	3.30 × 10 ⁸	3.20 × 10 ⁸	3.79 × 10 ⁸	2.19 × 10 ⁸	2.13 × 10 ⁸	- ^c	7.78 × 10 ⁷	7.34 × 10 ⁷	-	3.04 × 10 ⁸	3.03 × 10 ⁸	4.47 × 10 ⁸
	1.51 × 10 ⁸	1.47 × 10 ⁸	1.96 × 10 ⁸	8.89 × 10 ⁷	8.77 × 10 ⁷	-	6.56 × 10 ⁷	6.21 × 10 ⁷	-	1.19 × 10 ⁸	1.18 × 10 ⁸	2.87 × 10 ⁸
	1.18 × 10 ⁸	1.14 × 10 ⁸	1.50 × 10 ⁸	7.51 × 10 ⁷	7.37 × 10 ⁷	-	6.01 × 10 ⁷	5.68 × 10 ⁷	-	1.00 × 10 ⁸	9.99 × 10 ⁷	1.73 × 10 ⁸
<i>S. epidermidis</i>	3.43 × 10 ⁸	3.38 × 10 ⁸	3.43 × 10 ⁸	1.65 × 10 ⁷	1.50 × 10 ⁷	1.59 × 10 ⁸	3.06 × 10 ⁷	3.03 × 10 ⁷	-	1.75 × 10 ⁸	1.73 × 10 ⁸	3.96 × 10 ⁸
	1.73 × 10 ⁸	1.70 × 10 ⁸	1.59 × 10 ⁸	4.37 × 10 ⁷	3.66 × 10 ⁷	1.19 × 10 ⁸	6.91 × 10 ⁷	6.89 × 10 ⁷	-	1.57 × 10 ⁸	1.55 × 10 ⁸	1.59 × 10 ⁸
	8.41 × 10 ⁷	2.96 × 10 ⁷	6.67 × 10 ⁷	3.67 × 10 ⁷	2.94 × 10 ⁷	5.32 × 10 ⁷	5.34 × 10 ⁷	5.30 × 10 ⁷	-	7.56 × 10 ⁷	7.42 × 10 ⁷	7.96 × 10 ⁷
	4.10 × 10 ⁷	1.87 × 10 ⁷	2.69 × 10 ⁷	2.14 × 10 ⁷	1.63 × 10 ⁷	3.98 × 10 ⁷	2.88 × 10 ⁷	2.85 × 10 ⁷	-	3.57 × 10 ⁷	3.48 × 10 ⁷	2.69 × 10 ⁷
	4.04 × 10 ⁷	1.48 × 10 ⁷	1.37 × 10 ⁷	1.74 × 10 ⁷	1.32 × 10 ⁷	2.69 × 10 ⁷	3.27 × 10 ⁷	3.25 × 10 ⁷	0	3.99 × 10 ⁷	3.87 × 10 ⁷	2.69 × 10 ⁷
<i>E. faecalis</i>	2.33 × 10 ⁹	2.32 × 10 ⁹	2.33 × 10 ⁹	1.20 × 10 ⁹	1.16 × 10 ⁹	8.82 × 10 ⁸	5.54 × 10 ⁸	5.33 × 10 ⁸	-	7.10 × 10 ⁸	7.07 × 10 ⁸	2.02 × 10 ⁹
	1.20 × 10 ⁹	1.19 × 10 ⁹	1.27 × 10 ⁹	1.44 × 10 ⁸	1.26 × 10 ⁸	-	8.43 × 10 ⁶	8.10 × 10 ⁶	-	3.17 × 10 ⁸	3.14 × 10 ⁸	1.41 × 10 ⁹
	5.86 × 10 ⁸	5.68 × 10 ⁸	5.94 × 10 ⁸	4.26 × 10 ⁷	4.00 × 10 ⁷	-	4.30 × 10 ⁶	4.20 × 10 ⁶	-	1.43 × 10 ⁸	1.42 × 10 ⁸	5.94 × 10 ⁸
	2.78 × 10 ⁸	2.74 × 10 ⁸	2.60 × 10 ⁸	4.00 × 10 ⁷	3.87 × 10 ⁷	-	2.02 × 10 ⁷	1.98 × 10 ⁷	-	1.20 × 10 ⁸	1.17 × 10 ⁸	3.37 × 10 ⁸
	2.27 × 10 ⁸	2.21 × 10 ⁸	2.08 × 10 ⁸	3.62 × 10 ⁷	3.53 × 10 ⁷	-	2.52 × 10 ⁷	2.48 × 10 ⁷	-	1.16 × 10 ⁸	1.13 × 10 ⁸	2.86 × 10 ⁸
<i>E. coli</i>	6.04 × 10 ⁸	5.57 × 10 ⁸	6.04 × 10 ⁸	8.96 × 10 ⁷	7.17 × 10 ⁷	2.94 × 10 ⁸	1.69 × 10 ⁷	1.50 × 10 ⁷	-	2.17 × 10 ⁸	2.04 × 10 ⁸	5.51 × 10 ⁸
	2.98 × 10 ⁸	2.76 × 10 ⁸	3.21 × 10 ⁸	6.04 × 10 ⁷	4.17 × 10 ⁷	9.85 × 10 ⁷	4.89 × 10 ⁷	4.39 × 10 ⁷	-	2.07 × 10 ⁸	1.93 × 10 ⁸	3.38 × 10 ⁸
	1.51 × 10 ⁸	1.41 × 10 ⁸	1.52 × 10 ⁸	4.80 × 10 ⁷	3.42 × 10 ⁷	-	5.99 × 10 ⁷	5.11 × 10 ⁷	-	1.38 × 10 ⁸	1.23 × 10 ⁸	1.87 × 10 ⁸
	6.55 × 10 ⁷	6.02 × 10 ⁷	6.34 × 10 ⁷	3.75 × 10 ⁷	2.51 × 10 ⁷	-	5.12 × 10 ⁷	4.20 × 10 ⁷	-	6.31 × 10 ⁷	5.55 × 10 ⁷	8.11 × 10 ⁷
	5.47 × 10 ⁷	5.20 × 10 ⁷	3.68 × 10 ⁷	3.28 × 10 ⁷	1.87 × 10 ⁷	-	4.47 × 10 ⁷	4.07 × 10 ⁷	-	5.10 × 10 ⁷	4.44 × 10 ⁷	8.11 × 10 ⁷

^aBacterial cell number was measured by flow cytometry (FCM) and spectrophotometer method of optical density (OD) measurement after 1 hr exposure to ZnO, TiO₂ and SiO₂ nanoparticles; inoculum used for each experiment was indicated in the control samples, i.e. no nanoparticles.

^bPresented data were converted from each sample cells concentration according to the each species standard curve of cell/ml vs OD₆₆₀ and as mean of triplicate with standard deviations (SD) of < 5%.

^cValue was negative.

dispersed in Milli-Q water, and 3 μ L suspensions was deposited on the TEM grid, dried, and evacuated before analysis. Images were collected using a field emission JEM-2100 F (JEOL, Tokyo, Japan) equipped with a CCD camera in high resolution mode with an acceleration voltage of 100 kV.

The hydrodynamic size and zeta potential were measured in Milli-Q water using a Zetasizer (Malvern, Worcestershire, UK) as described in previous study [43]. Briefly, the nanoparticle samples were measured after dilution of a nanoparticle stock solution to 50 μ g/ml in Milli-Q water. These dilutions were sonicated for 30 min and vortexed briefly to provide a homogenous dispersion. For the size measurement, 70 μ L of the diluted dispersion nanoparticles was transferred to a cuvette for dynamic size measurement; for zeta potential measurement, a Malvern zeta potential cell was washed three times with ultrapure water followed by transferring 850 μ L of diluted dispersion nanoparticles to this cell to measure the zeta potential. The concentration of the samples and experimental methods were optimized to assure the quality of the data. NIST standard gold nanoparticles (10 nm, 30 nm, and 60 nm) were used in the validation of the instrument. Both size and zeta potential were measured at least three times. The data were calculated as the average size or zeta potential of nanoparticles.

Bacterial strains and culture conditions

Four bacterial species were chosen for all experiments (Table 2). The bacterial stock cultures were stored in freezer (-80°C) with glycerol to a final concentration of 15%. *E. faecalis* and *E. coli* from the glycerol stocks were streaked into brain heart infusion (BHI) agar plates at 37°C overnight in an anaerobic chamber (Coy Laboratory Products INC.). For *S. enterica* Newport and *S. epidermidis*, the plates were grown under aerobic condition. One colony was picked by a loop and inoculated into a 50-ml Falcon centrifuge tube containing 10 ml BHI medium. The cultures were incubated anaerobically or aerobically in static conditions at 37°C overnight for use as seed cultures. Each seed culture of the bacteria was inoculated into BHI medium with an inoculation ratio of 1% (v/v), except *S. epidermidis* with 10% (v/v) of inoculation into BHI medium, and then the cultures were incubated at 37°C under anaerobic or aerobic condition without agitation for 4–6 hrs to enter the exponential phase based on our preliminary experiments [40]. The cells were collected by centrifugation (10,000 rpm, 2 min, 4°C), washed twice and then re-suspended in sterile saline solution (0.85% NaCl), which served as experimental bacterial cells. To measure the bacterial numbers in the presence of different concentrations of nano ZnO, TiO₂, and SiO₂, various concentrations of the nanoparticles (final concentrations were 0, 0.1, 0.2, 0.3, 0.5, 1 mg/ml) based on our

preliminary experiments were added into each bacterial cell re-suspension and mixed well by vortexing, leaving one as a control without nanoparticles, but same volume of Milli-Q water, and then kept in the dark for 1 hr at 4°C [39]. In order to test the different bacterial concentrations after exposure to the nanoparticles, the initial bacterial re-suspension with approximately 10^8 - 10^9 cells/ml was serially diluted (0, -2, -4, -8, -10 fold) in saline solution and then mixed well with each nanoparticles at final concentration of 0.5 mg/ml, 0.5 mg/ml and 1 mg/ml for ZnO, TiO₂ and SiO₂, respectively. All the samples were kept under the same conditions as mentioned above. A control (containing saline solution and nanoparticles without bacterial cells) was included in all experiments and kept under same conditions. All experiments were carried out in triplicates. After 1 hr exposure to the nanoparticles, the bacterial cell concentrations were measured by different methods as mentioned below [40,41,44,45].

Plate counting cell numbers


Samples were withdrawn and then serially diluted in saline solution. Aliquots of 10 μ L were spread on BHI-plates. After overnight incubation at 37°C , colonies on the plates were counted to determine the number of CFU [46].

Optical density measurement

Aliquots of 200 μ L were withdrawn, added to a 96-well plate (Corning incorporated, flat bottom, non-lid) and immediately assayed by measuring the optical density in a SpectraMax M2 plate reader (Molecular Devices) at 660 nm [41]. The absorbance values of the controls were subtracted from the experimental values [36].

Flow cytometry analysis of bacterial cell numbers in combination with LIVE/DEAD BacLight bacterial viability and counting kit

Samples were collected, diluted and stained according to the manufacture's instruction using the BacLight LIVE/DEAD bacterial viability and counting kit as described briefly here. Each of 1.5 μ L of 3.34 mM SYTO 9 green fluorescent nucleic acid dye (Component A) and of 20 mM propidium iodide (Component B) was added to the flow cytometry tube containing 1 ml sample. Incubate the sample for 15 minutes at room temperature protected from light. The bacterial cell numbers assay was performed on the Accuri C6 flow cytometry. Fluorescence filters and detectors were all standardized with green fluorescence collected in the FL1 channel (530 ± 15 nm) and red fluorescence collected in the FL3 channel (>670 nm). All parameters were collected as logarithmic signals. A similar setup of parameters was used as described previously [40]. Data were analyzed using CFlow Plus software. In density plots of light scatter properties, bacterial cells were gated from irrelevant counts for



fluorescence analyses. In density plots of fluorescence, the distinct bacterial populations (live cells and damaged or dead cells) were gated based on the different viability stages. Total cell numbers = live cell numbers + dead cell numbers. Accuri C6 flow cytometry was calibrated using 8-peak Spherotech Validation Beads m.

Standard curve of optical density versus cell number for each bacterial strain

Exponentially growing cells of each bacterial species were serially diluted in saline solution in triplicate. Then OD₆₆₀ of the samples was measured by above mentioned method. Sterile saline solution was used as blanks. For counting cell numbers, the serially diluted bacterial cultures were further diluted to 1 ml with saline solution. Then the total bacterial cell number was analyzed by

flow cytometry as mentioned above. The correlation between OD₆₆₀ and cells number for each bacterial species was established by means of a standard curve (Figure 3).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

HP, YZ, GH, NK, and HC performed research and analyzed data. HC conceived and designed the project. HC wrote the paper with help from all authors. The final manuscript was read and approved by all authors.

Acknowledgements

We would like to thank Drs. Steven L. Foley and Jing Han for their critical review of this manuscript. This study was funded by National Center for Toxicological Research, United States Food and Drug Administration, and supported in part by appointment (H.P.) to the Postgraduate Research Fellowship Program by the Oak Ridge Institute for Science and Education through an interagency agreement between the U.S. Department of Energy and the U.S. Food and Drug Administration. The authors would like to thank

M. Yvonne Jones for assistance with TEM images. The views presented in this article do not necessarily reflect those of the Food and Drug Administration.

Author details

¹Division of Microbiology, National Center for Toxicological Research, U.S. Food and Drug Administration, Jefferson, AR 72079, USA. ²Nanotechnology Core Facility, National Center for Toxicological Research, U.S. Food and Drug Administration, Jefferson, AR 72079, USA. ³Department of Clinical Laboratory and Nutritional Sciences, University of Massachusetts Lowell, Lowell, MA 01854, USA.

Received: 17 March 2014 Accepted: 13 August 2014

Published: 21 August 2014

References

- Hajipour MJ, Fromm KM, Ashkarran AA, Jimenez de Aberasturi D, De Larramendi IR, Rojo T, Serpooshan V, Parak WJ, Mahmoudi M: **Antibacterial properties of nanoparticles.** *Trends Biotechnol* 2012, **30**(10):499–511.
- Allahverdiyev AM, Abamor ES, Bagirova M, Rafailovich M: **Antimicrobial effects of TiO₂ and Ag₂O nanoparticles against drug-resistant bacteria and leishmania parasites.** *Future Microbiol* 2011, **6**(8):933–940.
- Suresh AK, Pelletier DA, Doktycz MJ: **Relating nanomaterial properties and microbial toxicity.** *Nanoscale* 2013, **5**(2):463–474.
- Valdiglesias V, Costa C, Kilic G, Costa S, Pasaro E, Laffon B, Teixeira JP: **Neuronal cytotoxicity and genotoxicity induced by zinc oxide nanoparticles.** *Environ Int* 2013, **55**:92–100.
- Warheit DB: **How to measure hazards/risks following exposures to nanoscale or pigment-grade titanium dioxide particles.** *Toxicol Lett* 2013, **220**(2):193–204.
- Hoff D, Sheikh L, Bhattacharya S, Nayar S, Webster TJ: **Comparison study of ferrofluid and powder iron oxide nanoparticle permeability across the blood–brain barrier.** *Int J Nanomedicine* 2013, **8**:703–710.
- Thorley AJ, Tetley TD: **New perspectives in nanomedicine.** *Pharmacol Ther* 2013, **140**(2):176–185.
- Ko H-H, Chen H-T, Yen F-L, Lu W-C, Kuo C-W, Wang M-C: **Preparation of TiO₂ Nanocrystallite Powders Coated with 9 mol% ZnO for Cosmetic Applications in Sunscreens.** *Int J Mol Sci* 2012, **13**(2):1658–1669.
- Battez AH, Gonzalez R, Viesca JL, Fernandez JE, Fernandez JMD, Machado A, Chou R, Riba J: **CuO, ZrO₂ and ZnO nanoparticles as antiwear additive in oil lubricants.** *Wear* 2008, **265**(3–4):422–428.
- Duncan TV: **Applications of nanotechnology in food packaging and food safety: barrier materials, antimicrobials and sensors.** *J Colloid Interface Sci* 2011, **363**(1):1–24.
- Gupta S, Tripathi M: **A review of TiO₂ nanoparticles.** *Chin Sci Bull* 2011, **56**(16):1639–1657.
- Applerot G, Lipovsky A, Dror R, Perkas N, Nitzan Y, Lubart R, Gedanken A: **Enhanced antibacterial activity of nanocrystalline ZnO Due to increased ROS-mediated cell injury.** *Adv Funct Mater* 2009, **19**(6):842–852.
- Warnes SL, Caves V, Keevil CW: **Mechanism of copper surface toxicity in Escherichia coli O157:H7 and Salmonella involves immediate membrane depolarization followed by slower rate of DNA destruction which differs from that observed for Gram-positive bacteria.** *Environ Microbiol* 2012, **14**(7):1730–1743.
- Jena P, Mohanty S, Mallick R, Jacob B, Sonawane A: **Toxicity and antibacterial assessment of chitosan-coated silver nanoparticles on human pathogens and macrophage cells.** *Int J Nanomedicine* 2012, **7**:1805–1818.
- Wagenvoort JHT, De Brauwier EIGB, Penders RJR, Willems RJ, Top J, Bonten MJ: **Environmental survival of vancomycin-resistant Enterococcus faecium.** *J Hosp Infect* 2011, **77**(3):282–283.
- Seil JT, Webster TJ: **Antimicrobial applications of nanotechnology: methods and literature.** *Int J Nanomedicine* 2012, **7**:2767–2781.
- Saravanan M, Nanda A: **Extracellular synthesis of silver bionanoparticles from Aspergillus clavatus and its antimicrobial activity against MRSA and MRSE.** *Colloids Surf B: Biointerfaces* 2010, **77**(2):214–218.
- De Lima R, Seabra AB, Duran N: **Silver nanoparticles: a brief review of cytotoxicity and genotoxicity of chemically and biogenically synthesized nanoparticles.** *J Appl Toxicol* 2012, **32**(11):867–879.
- Bondarenko O, Juganson K, Ivask A, Kasemets K, Mortimer M, Kahru A: **Toxicity of Ag, CuO and ZnO nanoparticles to selected environmentally relevant test organisms and mammalian cells in vitro: a critical review.** *Arch Toxicol* 2013, **87**(7):1181–1200.
- Quigley L, O'Sullivan O, Beresford TP, Ross RP, Fitzgerald GF, Cotter PD: **Molecular approaches to analysing the microbial composition of raw milk and raw milk cheese.** *Int J Food Microbiol* 2011, **150**(2–3):81–94.
- Fang H, Xu J, Ding D, Jackson SA, Patel IR, Frye JG, Zou W, Nayak R, Foley S, Chen J, Su J, Ye Y, Turner S, Harris S, Zhou G, Cerniglia C, Tong W: **An FDA bioinformatics tool for microbial genomics research on molecular characterization of bacterial foodborne pathogens using microarrays.** *BMC Bioinformatics* 2010, **11**(Suppl 6):S4.
- Zhang K, Cheng L, Imazato S, Antonucci JM, Lin NJ, Lin-Gibson S, Bai Y, Xu HHK: **Effects of dual antibacterial agents MDPB and nano-silver in primer on microcosm biofilm, cytotoxicity and dentine bond properties.** *J Dent* 2013, **41**(5):464–474.
- Koseki S, Nonaka J: **Alternative approach to modeling bacterial lag time, using logistic regression as a function of time, temperature, pH, and sodium chloride concentration.** *Appl Environ Microbiol* 2012, **78**(17):6103–6112.
- Schacht VJ, Neumann LV, Sandhi SK, Chen L, Henning T, Klar PJ, Theophel K, Schnell S, Bunge M: **Effects of silver nanoparticles on microbial growth dynamics.** *J Appl Microbiol* 2013, **114**(1):25–35.
- Dudak FC, Boyaci IH: **Rapid and label-free bacteria detection by surface plasmon resonance (SPR) biosensors.** *Biotechnol J* 2009, **4**(7):1003–1011.
- Vital M, Dignum M, Magic-Knezev A, Ross P, Rietveld L, Hammes F: **Flow cytometry and adenosine tri-phosphate analysis: alternative possibilities to evaluate major bacteriological changes in drinking water treatment and distribution systems.** *Water Res* 2012, **46**(15):4665–4676.
- Zahavy E, Ber R, Gur D, Abramovich H, Freeman E, Maoz S, Yitzhaki S: **Application of nanoparticles for the detection and sorting of pathogenic bacteria by flow-cytometry.** *Adv Exp Med Biol* 2012, **733**:23–36.
- Masco L, Vanhoutte T, Temmerman R, Swings J, Huys G: **Evaluation of real-time PCR targeting the 16S rRNA and recA genes for the enumeration of bifidobacteria in probiotic products.** *Int J Food Microbiol* 2007, **113**(3):351–357.
- Lazcka O, Del Campo FJ, Munoz FX: **Pathogen detection: a perspective of traditional methods and biosensors.** *Biosens Bioelectron* 2007, **22**(7):1205–1217.
- Davey HM: **Life, death, and in-between: meanings and methods in microbiology.** *Appl Environ Microbiol* 2011, **77**(16):5571–5576.
- Kerstens M, Boulet G, Pintelon I, Hellings M, Voeten L, Delputte P, Maes L, Cos P: **Quantification of Candida albicans by flow cytometry using TO-PRO[®](-3) iodide as a single-stain viability dye.** *J Microbiol Methods* 2013, **92**(2):189–191.
- Lehtinen J, Nuutila J, Lilius E-M: **Green fluorescent protein-propidium iodide (GFP-PI) based assay for flow cytometric measurement of bacterial viability.** *Cytometry A* 2004, **60**(2):165–172.
- Hammes F, Egli T: **Cytometric methods for measuring bacteria in water: advantages, pitfalls and applications.** *Anal Bioanal Chem* 2010, **397**(3):1083–1095.
- Muller S, Nebe-von-Caron G: **Functional single-cell analyses: flow cytometry and cell sorting of microbial populations and communities.** *FEMS Microbiol Rev* 2010, **34**(4):554–587.
- Mallick S, Sharma S, Banerjee M, Ghosh SS, Chattopadhyay A, Paul A: **Iodine-stabilized Cu nanoparticle chitosan composite for antibacterial applications.** *ACS Appl Mater Interfaces* 2012, **4**(3):1313–1323.
- Sadiq IM, Chandrasekaran N, Mukherjee A: **Studies on Effect of TiO₂ Nanoparticles on Growth and Membrane Permeability of Escherichia coli, Pseudomonas aeruginosa, and Bacillus subtilis.** *Curr Nanosci* 2010, **6**(4):381–387.
- Padmavathy N, Vijayaraghavan R: **Interaction of ZnO nanoparticles with microbes-a physio and biochemical assay.** *J Biomed Nanotechnol* 2011, **7**(6):813–822.
- Fang T-T, Li X, Wang Q-S, Zhang Z-J, Liu P, Zhang C-C: **Toxicity evaluation of CdTe quantum dots with different size on Escherichia coli.** *Toxicol In Vitro* 2012, **26**(7):1233–1239.
- Kumar A, Pandey AK, Singh SS, Shanker R, Dhawan A: **Engineered ZnO and TiO₂ nanoparticles induce oxidative stress and DNA damage leading to reduced viability of Escherichia coli.** *Free Radic Biol Med* 2011, **51**(10):1872–1881.
- Pan H, Feng J, Cerniglia CE, Chen H: **Effects of Orange II and Sudan III azo dyes and their metabolites on Staphylococcus aureus.** *J Ind Microbiol Biotechnol* 2011, **38**(10):1729–1738.

41. Pan H, Feng J, He G-X, Cerniglia CE, Chen H: **Evaluation of impact of exposure of Sudan azo dyes and their metabolites on human intestinal bacteria.** *Anaerobe* 2012, **18**(4):445–453.
42. Sharma V, Shukla RK, Saxena N, Parmar D, Das M, Dhawan A: **DNA damaging potential of zinc oxide nanoparticles in human epidermal cells.** *Toxicol Lett* 2009, **185**(3):211–218.
43. Zhang Y, Ferguson SA, Watanabe F, Jones Y, Xu Y, Biris AS, Hussain S, Ali SF: **Silver nanoparticles decrease body weight and locomotor activity in adult male rats.** *Small* 2013, **9**(9–10):1715–1720.
44. Xu H, Heinze TM, Paine DD, Cerniglia CE, Chen H: **Sudan azo dyes and Para Red degradation by prevalent bacteria of the human gastrointestinal tract.** *Anaerobe* 2010, **16**(2):114–119.
45. Stingley RL, Zou W, Heinze TM, Chen H, Cerniglia CE: **Metabolism of azo dyes by human skin microbiota.** *J Med Microbiol* 2010, **59**(Pt 1):108–114.
46. Mangoni ML, Papo N, Barra D, Simmaco M, Bozzi A, Di Giulio A, Rinaldi AC: **Effects of the antimicrobial peptide temporin L on cell morphology, membrane permeability and viability of Escherichia coli.** *Biochem J* 2004, **380**(Pt 3):859–865.

doi:10.1186/s12866-014-0222-6

Cite this article as: Pan *et al.*: A comparison of conventional methods for the quantification of bacterial cells after exposure to metal oxide nanoparticles. *BMC Microbiology* 2014 **14**:222.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

